

Neurofibromin (McNFn27a): sc-20017

BACKGROUND

Neurofibromatosis type 1 (NF1), or von Recklinghausen neurofibromatosis, is one of the most common autosomal dominant disorders in humans. Early linkage analysis mapped the NF1 gene to chromosome 17q11.2. The predicted NF1 transcript encodes a 2,818 amino acid protein designated NF1GRP. By sequence analysis, similarity has been demonstrated within a small region of NF1GRP and members of the Ras GAP gene family. Functionally, NF1GRP was shown by biochemical analysis involving RAS-GAP hydrolysis and functional complementation in yeast to further resemble GAP protein. The NF1 protein is expressed at relatively constant levels in a broad range of cell lines and tissues including brain, lung, liver, kidney, spleen, muscle and colon. Although little is known regarding the function of NF1GRP, the homology with the catalytic domain of proteins with GTPase activity suggests that the NF1GRP may also interact *in vivo* with Ras proteins.

CHROMOSOMAL LOCATION

Genetic locus: NF1 (human) mapping to 17q11.2; Nf1 (mouse) mapping to 11 B5.

SOURCE

Neurofibromin (McNFn27a) is a mouse monoclonal antibody raised against amino acids 27-41 corresponding to an N-terminal peptide of Neurofibromin of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Neurofibromin (McNFn27a) is recommended for detection of Neurofibromin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Neurofibromin siRNA (h): sc-36036, Neurofibromin siRNA (m): sc-36037, Neurofibromin shRNA Plasmid (h): sc-36036-SH, Neurofibromin shRNA Plasmid (m): sc-36037-SH, Neurofibromin shRNA (h) Lentiviral Particles: sc-36036-V and Neurofibromin shRNA (m) Lentiviral Particles: sc-36037-V.


Molecular Weight of Neurofibromin: 250 kDa.

Positive Controls: H4 cell lysate: sc-2408, A-431 whole cell lysate: sc-2201 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Neurofibromin (McNFn27a): sc-20017. Western blot analysis of Neurofibromin expression in NF1 knockout (A), rat (B) and human (C) Schwann cells. Kindly provided by Dr. David Muir, University of Florida College of Medicine.



Neurofibromin (McNFn27a): sc-20017. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing cytoplasmic staining of hepatic sinusoids.

SELECT PRODUCT CITATIONS

- Caudell, D., et al. 2010. Retroviral insertional mutagenesis identifies Zeb2 activation as a novel leukemogenic collaborating event in CALM-AF10 transgenic mice. *Blood* 115: 1194-1203.
- Godin, F., et al. 2012. A fraction of Neurofibromin interacts with PML bodies in the nucleus of the CCF astrocytoma cell line. *Biochem. Biophys. Res. Commun.* 418: 689-694.
- Kahen, E.J., et al. 2018. Neurofibromin level directs Ras pathway signaling and mediates sensitivity to targeted agents in malignant peripheral nerve sheath tumors. *Oncotarget* 9: 22571-22585.
- Ramakrishnan, G., et al. 2022. NF1 loss of function as an alternative initiating event in pancreatic ductal adenocarcinoma. *Cell Rep.* 41: 111623.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Neurofibromin (H-12): sc-376886** for Neurofibromin antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.