

Rad53 (γ-300): sc-20169

BACKGROUND

DNA damage results in the arrest of cell cycle progression, allowing the damaged DNA to be repaired prior to replication. Checkpoints exist at several cell cycle phase transitions to maintain this genetic integrity. Rad9, Rad17, Rad24 and Mec3 are involved in activating the G₁ and G₂ checkpoints. Pol2 (also known as Dun2), encoding the catalytic subunit of DNA polymerase ε, plays a role in activating the S phase checkpoint. The protein kinase Rad53 (also designated Spk1, Mec2 or Sad1) is essential for both G₂ and S phase arrest. Activation of Rad53 is regulated by Mec1 (also known as Esr1 and Sad3), a homolog of the human ATM protein. Pds1 and Mad2 both regulate checkpoints associated with incomplete spindle replication. Dun1, another protein kinase, plays a role in transducing the DNA damage signal.

REFERENCES

- Li, R., et al. 1993. The mitotic feedback control gene MAD2 encodes the α subunit of a prenyltransferase. *Nature* 366: 82-84.
- Zhou, Z. and Elledge, S.J. 1993. Dun1 encodes a protein kinase that controls the DNA damage response in yeast. *Cell* 75: 1119-1127.
- Abloussekhra, A., et al. 1996. A novel role for the budding yeast RAD9 checkpoint gene in DNA damage-dependent transcription. *EMBO J.* 15: 3912-3922.
- Siede, W., et al. 1996. Cloning and characterization of RAD17, a gene controlling cell cycle responses to DNA damage in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 24: 1669-1675.
- Lydall, D., et al. 1996. A meiotic recombination checkpoint controlled by mitotic checkpoint genes. *Nature* 383: 840-843.
- Longhese, M.P., et al. 1996. Yeast pep3/Mec3 mutants fail to delay entry into S phase and to slow DNA replication in response to DNA damage, and they define a functional link between Mec3 and DNA primase. *Mol. Cell Biol.* 16: 3235-3244.
- Navas, T.A., et al. 1996. Rad9 and DNA polymerase ε form parallel sensory branches for transducing the DNA damage checkpoint signal in *Saccharomyces cerevisiae*. *Genes Dev.* 10: 2632-2643.
- Yamamoto, A., et al. 1996. Pds1p, an inhibitor of anaphase in budding yeast, plays a critical role in the APC and checkpoint pathway(s). *J. Cell Biol.* 133: 99-110.

SOURCE

Rad53 (γ-300) is a rabbit polyclonal antibody raised against amino acids 1-300 of Rad53 of *Saccharomyces cerevisiae* origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS


Rad53 (γ-300) is recommended for detection of Rad53 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Rad53: 92 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



Rad53 (γ-300): sc-20169. Western blot analysis of yeast recombinant Rad53 fusion protein.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **Rad53 (A-9): sc-74427** or **Rad53 (B-6): sc-74426**, our highly recommended monoclonal alternatives to Rad53 (γ-300).