

normal goat IgG: sc-2028

BACKGROUND

Santa Cruz Biotechnology offers a wide variety of control immunoglobulin and control sera for a large selection of species, including mouse, rabbit, goat, chicken, rat, hamster, canine, guinea pig and sheep. Control immunoglobulin and immunoglobulin conjugates are useful negative controls. Normal sera is offered to be used as blocking reagents. Santa Cruz Biotechnology offers affinity purified normal immunoglobulins and immunoglobulin conjugates for use as negative controls in applications including flow cytometry, immunohistochemistry and immunofluorescence. Isotype specific control immunoglobulins include classes such as mouse IgG₁, IgG_{2a}, IgG_{2b}, IgG₃, IgM and IgA, rat IgG₁, IgG_{2a}, IgG_{2b} and IgM, Armenian hamster IgG, and both goat and rabbit IgG. Most control immunoglobulins are available as unconjugated controls or as FITC (fluorescein isothiocyanate), PE (phycoerythrin), PE-Cy5 (phycoerythrin-Cy5), PE-Cy7 (phycoerythrin-Cy7), APC (allophycocyanin) and APC-Cy7 (allophycocyanin-Cy7) conjugates. Additional conjugates include Alexa Fluor[®] 488, Alexa Fluor[®] 647, Alexa Fluor[®] 405, PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex-Cy 5.5).

SOURCE

normal goat IgG is an unconjugated, affinity purified isotype control immunoglobulin from goat.

PRODUCT

Each vial contains 200 µg goat IgG in 0.5 ml PBS with 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

normal goat IgG is recommended for use as a isotype control immunoglobulin in place of a target specific primary antibody of the same isotype (goat IgG) by Western blotting, immunofluorescence staining, immunohistochemical staining and flow cytometry. To be used at an assay dependent dilution.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

SELECT PRODUCT CITATIONS

- Ali, A.S., et al. 2000. Detection of hTERT protein by flow cytometry. *Leukemia* 14: 2176-2181.
- Skildum, A.J., et al. 2002. The cyclin-dependent kinase inhibitor p21WAF1/Cip1 is an antiestrogen-regulated inhibitor of Cdk4 in human breast cancer cells. *J. Biol. Chem.* 277: 5145-5152.
- Donev, R., et al. 2003. Recruitment of heterogeneous nuclear ribonucleoprotein A1 *in vivo* to the LMP/TAP region of the major histocompatibility complex. *J. Biol. Chem.* 278: 5214-5226.
- Iwahori, T. 2003. CYP3A4 inducible model for *in vitro* analysis of human drug metabolism using a bioartificial liver. *Hepatology* 37: 665-673.
- Schroder, M., et al. 2004. The unfolded protein response represses differentiation through the RPD3-SIN3 histone deacetylase. *EMBO J.* 23: 2281-2292.
- Kam, K.Y., et al. 2005. Oct-1 and nuclear factor Y bind to the SURG-1 element to direct basal and gonadotropin-releasing hormone (GnRH)-stimulated mouse GnRH receptor gene transcription. *Mol. Endoc.* 19: 148-162.
- Morales, P. and Reyes, P. 2006. Infection of human fallopian tube epithelial cells with *Neisseria gonorrhoeae* protects cells from tumor necrosis factor α -induced apoptosis. *Infect. Immun.* 74: 3643-3650.
- Deschênes, J. and Bourdeau, V. 2007. Regulation of GREB1 transcription by estrogen receptor α through a multipartite enhancer spread over 20 kb of upstream flanking sequences. *J. Biol. Chem.* 282: 17335-17339.
- Lampe, X., et al. 2008. An ultraconserved Hox-Pbx responsive element resides in the coding sequence of Hoxa2 and is active in rhombomere 4. *Nucleic Acids Res.* 36: 3214-3225.
- Maarof, G., et al. 2010. Interleukin-24 inhibits the plasma cell differentiation program in human germinal center B cells. *Blood.* 115: 1718-1726.
- Delfino, D.V., et al. 2011. Glucocorticoid-induced activation of caspase-8 protects the glucocorticoid-induced protein Gilz from proteasomal degradation and induces its binding to SUMO-1 in murine thymocytes. *Cell Death Differ.* 18: 183-190.
- Lindner, H.B. et al. 2011. Anti-bacterial effects of poly-N-acetyl-glucosamine nanofibers in cutaneous wound healing: requirement for Akt1. *PLoS ONE* 6: e18996.
- Li, B., et al. 2011. Increased Hedgehog signaling in postnatal kidney results in aberrant activation of nephron developmental programs. *Hum. Mol. Genet.* 20: 4155-4166.
- Aporntewan, C., et al. 2011. Hypomethylation of intragenic LINE-1 represses transcription in cancer cells through AGO2. *PLoS ONE* 6: e17934.
- Li, ZY., et al. 2011. Positive regulation of hepatic miR-122 expression by HNF4 α . *J. Hepatol.* 55: 602-611.
- Huang, L., et al. 2011. Prevention of transcriptional silencing by a replicator-binding complex consisting of SWI/SNF, MeCP1, and hnRNP C1/C2. *Mol. Cell. Biol.* 31: 3472-3484.