

donkey anti-goat IgG-B: sc-2042

BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. Secondary antibodies are commonly affinity purified against immobilized whole IgG or against antibody fragments. Santa Cruz Biotechnology offers an extensive selection of secondary antibodies optimized for immunohistochemistry and flow cytometry, and are labeled with either biotin, FITC (fluorescein isothiocyanate), Texas Red[®], TRITC (tetramethyl rhodamine isothiocyanate), PE (phycoerythrin), PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex with cyanin-5.5). Immunohistochemistry and flow cytometry secondary antibodies are specific for commonly used primary antibody species, including goat, rabbit, mouse and rat.

SOURCE

donkey anti-goat IgG-B is a pre-adsorbed, affinity purified secondary antibody raised in donkey against goat IgG and conjugated to biotin.

PRODUCT

Each vial contains 200 µg goat IgG (mouse and human adsorbed IgG) in 0.5 ml of PBS containing 0.02% sodium azide.

APPLICATIONS

donkey anti-goat IgG-B is recommended for detection of goat IgG by immunofluorescence staining (starting dilution: 1:100, dilution range: 1:100-1:400) and immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400).

RECOMMENDED SUPPORT PRODUCTS

A. TISSUE CULTURE CELLS

- CrystalCruz[™] Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- CrystalCruz[™] Micro Slides 75 x 25 mm; 72 frosted sides: sc-24976
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336

B. FROZEN TISSUE SECTIONS

- Organo/Limonene Mount, non-toxic alternative to Permount, 100 ml: sc-45087
- UltraCruz[™] Mounting Medium, aqueous-based, 10 ml: sc-24941
- ImmunoHistoMount, aqueous-based mounting medium, 30 ml: sc-45086
- Immuno In Situ Mount, for use with in situ hybridization, 30 ml: sc-45088

C. FORMALIN-FIXED, PARAFFIN-EMBEDDED TISSUE SECTIONS

- Paraffin, for the preparation of tissue samples for staining, 500 g: sc-286633
- Xylenes, mixed isomers with ethylbenzene, 500 ml: sc-237422
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973


RESEARCH USE

For research use only, not for use in diagnostic procedures.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



donkey anti-goat IgG-B: sc-2042. Immunoperoxidase staining of formalin-fixed, paraffin-embedded normal human skin showing localization within the cytoplasm. Antibody tested: Villin (C-19): sc-7672.



donkey anti-goat IgG-B: sc-2042. Immunoperoxidase staining of formalin fixed, paraffin-embedded mouse kidney tissue showing membrane and cytoplasmic localization. Antibody tested: NDRG1 (N-19): sc-19464.

SELECT PRODUCT CITATIONS

- Carlotto, F., et al. 1999. Activation of nuclear factor κ B in single living cells. Dependence of nuclear translocation and anti-apoptotic function on EGFPRELA concentration. J. Biol. Chem. 274: 37941-37949.
- Parkinson, N., et al. 2006. Mutation at the Evi1 locus in Junbo mice causes susceptibility to otitis media. PLoS Genet. 2: e149.
- Lam, N.P., et al. 2007. Age-dependent increase of discoidin domain receptor 2 and matrix metalloproteinase 13 expression in temporomandibular joint cartilage of type IX and type XI collagen-deficient mice. Arch. Oral Biol. 52: 579-584.
- Rached, E., et al. 2008. Evaluation of putative biomarkers of nephrotoxicity after exposure to ochratoxin A *in vivo* and *in vitro*. Toxicol. Sci. 103: 371-381.
- Varayoud, J., et al. 2008. Developmental exposure to Bisphenol A impairs the uterine response to ovarian steroids in the adult. Endocrinology 149: 5848-5860.
- Sieber, M., et al. 2009. Comparative analysis of novel noninvasive renal biomarkers and metabolomic changes in a rat model of gentamicin nephrotoxicity. Toxicol. Sci. 109: 336-349.
- Girasol, A., et al. 2009. Fyn mediates leptin actions in the thymus of rodents. PLoS ONE 4: e7707.
- Abdeen, A., et al. 2009. Correlation between clinical outcome and growth factor pathway expression in osteogenic sarcoma. Cancer 115: 5243-5250.
- North, M.L., et al. 2011. Augmentation of arginase 1 expression by exposure to air pollution exacerbates the airways hyperresponsiveness in murine models of asthma. Respir. Res. 12: 19.
- Daane, J.M., et al. 2011. Mesothelium of the murine allantois exhibits distinct regional properties. J. Morphol. 272: 536-556.

Texas Red[®] is a registered trademark of Molecular Probes (6/02).