



## CPTI-L (C-14): sc-20516

### BACKGROUND

The mitochondrial beta-oxidation of long-chain fatty acids is initiated by the sequential action of carnitine palmitoyltransferase (CPT) I (outer membrane and detergent labile) and II (inner membrane and detergent stable), together with carnitine carrier. CPTI catalyzes the first reaction in the transport of long-chain fatty acids from the cytoplasm to the mitochondrion, a rate-limiting step in beta-oxidation. Two types of CPTI are known, the muscle and liver isoforms. The human liver CPTI gene is located to chromosome 11q and the muscle type protein is specially expressed in heart and skeletal muscle. Membrane-bound CPTI, but not CPTII, is inhibited reversibly by malonyl-coenzyme A (CoA). CPTI is about 86 kDa in non-hepatic tissues and approximately 90–94 kDa in liver, depending upon species. For CPTII, it is about 70 kDa in rat tissues and about 68 kDa in all mouse tissues and human liver. Unlike CPTII, CPTI requires membrane integrity for catalytic function. In addition, glutamic acid 3 and histidine 5 are necessary for malonyl CoA inhibition and binding to liver CPTI, but not catalytic activity.

### REFERENCES

1. Pande, S.V., et al. 1976. Characterization of carnitine acylcarnitine translocase system of heart mitochondria. *J. Biol. Chem.* 251: 6683-6691.
2. McGarry, J.D., et al. 1989. Regulation of ketogenesis and the renaissance of carnitine palmitoyltransferase. *Diabetes/Metab. Rev.* 5: 271-284.
3. Woeltje, K.F., et al. 1990. Inter-tissue and inter-species characteristics of the mitochondrial carnitine palmitoyltransferase enzyme system. *J. Biol. Chem.* 265: 10714-10719.
4. Britton, C.H., et al. 1995. Human liver mitochondrial carnitine palmitoyltransferase I: characterization of its cDNA and chromosomal localization and partial analysis of the gene. *Proc. Natl. Acad. Sci. USA* 92: 1984-1988.
5. Yamazaki, N., et al. 1996. Isolation and characterization of cDNA and genomic clones encoding human muscle type carnitine palmitoyltransferase I. *Biochim. Biophys. Acta* 1307: 157-161.
6. Zhu, H., et al. 1997. Functional studies of yeast-expressed human heart muscle carnitine palmitoyltransferase I. *Arch. Biochem. Biophys.* 347: 53-61.
7. Yamazaki, N., et al. 1997. Structural features of the gene encoding human muscle type carnitine palmitoyltransferase I. *FEBS Letts.* 409: 401-406.
8. Woldegiorgis, G., et al. 2000. Functional characterization of mammalian mitochondrial carnitine palmitoyltransferases I and II expressed in the yeast *Pichia pastoris*. *J. Nutr.* 130: 310-314.

### SOURCE

CPTI-L (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of CPTI-L of human origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-20516 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

CPTI-L (C-14) is recommended for detection of CPTI-L of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CPTI-L siRNA (h): sc-40380.

Molecular Weight of CPTI-L: 90-94 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.