

CPTII (S-15): sc-20526

BACKGROUND

The mitochondrial β -oxidation of long-chain fatty acids is initiated by the sequential action of carnitine palmitoyltransferase (CPT) I (outer membrane and detergent labile) and II (inner membrane and detergent stable), together with carnitine carrier. CPTI catalyzes the first reaction in the transport of long-chain fatty acids from the cytoplasm to the mitochondrion, a rate-limiting step in β -oxidation. Two types of CPT are known, the liver (CPTIA) and muscle (CPTIB) isoforms. The muscle type protein is specially expressed in heart and skeletal muscle. Membrane-bound CPTI, but not CPTII, is inhibited reversibly by malonyl-coenzyme A (CoA). Unlike CPTII, CPTI requires membrane integrity for catalytic function. In addition, glutamic acid 3 and histidine 5 are necessary for malonyl CoA inhibition and binding to liver CPTI, but not for catalytic activity.

REFERENCES

1. Pande, S.V., et al. 1976. Characterization of carnitine acylcarnitine translocase system of heart mitochondria. *J. Biol. Chem.* 251: 6683-6691.
2. McGarry, J.D., et al. 1989. Regulation of ketogenesis and the renaissance of carnitine palmitoyltransferase. *Diabetes Metab. Rev.* 5: 271-284.
3. Woeltje, K.F., et al. 1990. Inter-tissue and inter-species characteristics of the mitochondrial carnitine palmitoyltransferase enzyme system. *J. Biol. Chem.* 265: 10714-10719.
4. Britton, C.H., et al. 1995. Human liver mitochondrial carnitine palmitoyltransferase I: characterization of its cDNA and chromosomal localization and partial analysis of the gene. *Proc. Natl. Acad. Sci. USA* 92: 1984-1988.
5. Yamazaki, N., et al. 1996. Isolation and characterization of cDNA and genomic clones encoding human muscle type carnitine palmitoyltransferase I. *Biochim. Biophys. Acta* 1307: 157-161.
6. Zhu, H., et al. 1997. Functional studies of yeast-expressed human heart muscle carnitine palmitoyltransferase I. *Arch. Biochem. Biophys.* 347: 53-61.
7. Yamazaki, N., et al. 1997. Structural features of the gene encoding human muscle type carnitine palmitoyltransferase I. *FEBS Lett.* 409: 401-406.
8. Woldegiorgis, G., et al. 2000. Functional characterization of mammalian mitochondrial carnitine palmitoyltransferases I and II expressed in the yeast *Pichia pastoris*. *J. Nutr.* 130: 310-314.

CHROMOSOMAL LOCATION

Genetic locus: CPT2 (human) mapping to 1p32.3; Cpt2 (mouse) mapping to 4 C7.

SOURCE

CPTII (S-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CPTII of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-20526 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CPTII (S-15) is recommended for detection of CPTII of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CPTII (S-15) is also recommended for detection of CPTII in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for CPTII siRNA (h): sc-40378, CPTII siRNA (m): sc-40379, CPTII shRNA Plasmid (h): sc-40378-SH, CPTII shRNA Plasmid (m): sc-40379-SH, CPTII shRNA (h) Lentiviral Particles: sc-40378-V and CPTII shRNA (m) Lentiviral Particles: sc-40379-V.

Molecular Weight of CPTII: 67 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227 or mouse liver extract: sc-2256.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **CPTII (G-5): sc-377294**, our highly recommended monoclonal alternative to CPTII (S-15).