

band 3 (C-17): sc-20559

BACKGROUND

Band 3, also designated AE1, is an erythrocyte membrane glycoprotein that contributes to cell structural integrity and mediates exchange of chloride and bicarbonate across the phospholipid bilayer. The diverse functions of the approximately 900 amino acid protein are mediated by two distinct domains. The amino terminal domain, also known as cdb3 for cytoplasmic domain of erythrocyte membrane band 3, acts as an attachment site for the erythrocyte skeleton by binding ankyrin. The carboxy-terminal, membrane-associated domain carries out exchange transport of anions. Degradation of band 3 can generate an aging antigen known as senescent cell antigen, or SCA, which is expressed on old cells and marks them for removal by the immune system. An isoform of band 3, which lacks the first 65 amino acids and does not bind ankyrin, is expressed in kidney.

REFERENCES

1. Brock, C.J., et al. 1983. The human erythrocyte anion-transport protein. Partial amino acid sequence, conformation and a possible molecular mechanism for anion exchange. *Biochem. J.* 213: 577-586.
2. Kollert-Jons, A., et al. 1993. Anion exchanger 1 in human kidney and oncocyoma differs from erythroid AE1 in its NH₂ terminus. *Am. J. Physiol.* 265: F813-F821.
3. Jay, D.G. 1996. Role of band 3 in homeostasis and cell shape. *Cell* 86: 853-854.
4. Motais, R., et al. 1997. Association of the band 3 protein with a volume-activated, anion and amino acid channel: a molecular approach. *J. Exp. Biol.* 200: 361-367.
5. Tanner, M.J. 1997. The structure and function of band 3(AE1): recent developments (review). *Mol. Membr. Biol.* 14: 155-165.
6. Online Mendelian Inheritance in Man, OMIM™ 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 109270. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Mandal, D., et al. 2003. Caspase 3-mediated proteolysis of the N-terminal cytoplasmic domain of the human erythroid anion exchanger 1 (band 3). *J. Biol. Chem.* 278: 52551-52558.
8. Fu, G., et al. 2004. Purification and characterization of the human erythrocyte band 3 protein C-terminal domain. *Biochemistry* 43: 1633-1638.

CHROMOSOMAL LOCATION

Genetic locus: SLC4A1 (human) mapping to 17q21.31; Slc4a1 (mouse) mapping to 11 D.

SOURCE

band 3 (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of band 3 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-20559 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

band 3 (C-17) is recommended for detection of band 3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

band 3 (C-17) is also recommended for detection of band 3 in additional species, including porcine.

Suitable for use as control antibody for band 3 siRNA (h): sc-42735, band 3 siRNA (m): sc-42736, band 3 shRNA Plasmid (h): sc-42735-SH, band 3 shRNA Plasmid (m): sc-42736-SH, band 3 shRNA (h) Lentiviral Particles: sc-42735-V and band 3 shRNA (m) Lentiviral Particles: sc-42736-V.

Molecular Weight of band 3: 95 kDa.

Positive Controls: Caki-1 cell lysate: sc-2224 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **band 3 (A-6): sc-133190** or **band 3 (Bill 136): sc-58695**, our highly recommended monoclonal alternatives to band 3 (C-17).