

donkey anti-goat IgG-HRP: sc-2056

BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. Secondary antibodies are commonly affinity purified against immobilized whole IgG isotypes, including IgG₁, IgG_{2a}, IgG_{2b}, IgG₃ and IgG₄. Santa Cruz Biotechnology offers a wide selection of secondary antibodies, which are used in conjunction with our Cruz Marker™ molecular weight standards. We also provide specialized secondaries, such as pre-adsorbed secondary antibodies, which are pre-adsorbed with human IgG and mouse IgG for immunoglobulin-rich tissues and cells, F(ab')₂ fragment secondary antibodies that reduce non-specific secondary antibody binding to Fc receptors on the cell surface, and isotype-specific secondary antibodies against IgM, IgA and IgY.

SOURCE

donkey anti-goat IgG-HRP is a pre-adsorbed, affinity purified secondary antibody raised in donkey against goat IgG and conjugated to HRP (horseradish peroxidase).

PRODUCT

Each vial contains 200 µg mouse and human adsorbed IgG in 0.5 ml of 1X PBS containing 40% glycerol.

APPLICATIONS

donkey anti-goat IgG-HRP is recommended for detection of goat IgG by Western Blotting of immunoglobulin-rich tissues and cells (starting dilution: 1:5000, dilution range 1:5000-1:10000).

RECOMMENDED SUPPORT PRODUCTS

- Western Blotting Luminol Reagent, for 2,000 cm² membrane area: sc-2048
- RIPA Lysis Buffer, 50 ml, cell lysis buffer with protease inhibitors: sc-24948
- Electrophoresis Sample Buffer, 2X, 25 ml, reducing buffer: sc-24945
- Complete™ Protease Inhibitor Cocktail Tablet, 20 tablets: sc-29130
- Running Buffer, 10X, 1 L, TRIS-Glycine WB running buffer, pH 8.3: sc-24949
- Towbin, with SDS, 10X, 1 L, WB transfer buffer pH 8.3: sc-24954
- Bovine Serum Albumin (BSA), 100 g, blocking/incubation agent: sc-2323
- TBS Blotto A, lyophilized powder in single-use bottle: sc-2333
- UltraCruz™ PVDF Transfer Membrane, 0.45 µm, 30 cm x 3 m roll: sc-3723
- UltraCruz™ Nitrocellulose Pure Transfer Membrane, 0.22 µm, 30 cm x 3 m roll: sc-3718
- UltraCruz™ Electrophoresis Cell: sc-201625 : runs up to 10 or 15 sample by SDS – PAGE protein electrophoresis
- UltraCruz™ Autoradiography Film, Blue, 8 x 1, 100 sheets: sc-201697
- UltraCruz™ Gel Incubation Trays, 100 per pack: sc-201755 (blue), sc-201756 (green), sc-201757 (pink), sc-201758 (yellow), sc-201759 (orange)


STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



donkey anti-goat IgG-HRP: sc-2056. Western blot analysis of CALHM2 expression in HeLa (A), SH-SY5Y (B), SK-N-MC (C) and Hep G2 (D) whole cell lysates. Antibody tested: CALHM2 (E-12): sc-167376.



donkey anti-goat IgG-HRP: sc-2056. Western blot analysis of ABHD14A expression in HEL 92.1.7 whole cell lysate. Antibody tested: ABHD14A (T-14): sc-161301.

SELECT PRODUCT CITATIONS

- Schmitz, I., et al. 2003. Gene expression analysis of thymocyte selection *in vivo*. *Int. Immunol.* 15: 1237-1248.
- Popa-Nita, O., et al. 2008. Crystal-induced neutrophil activation: X. Pro-inflammatory role of the tyrosine kinase Tec. *Arthritis Rheum.* 58: 1866-1876.
- Rajamohan, S.B., et al. 2009. SIRT1 promotes cell survival under stress by deacetylation-dependent deactivation of poly (ADP-ribose) polymerase 1. *Mol. Cell. Biol.* 29: 4116-4129.
- Jayadev, S., et al. 2010. Presenilin 2 is the predominant γ -secretase in microglia and modulates cytokine release. *PLoS ONE* 5: e15743.
- Bohr, W., et al. 2010. Recombinant expression, purification, and functional characterisation of connective tissue growth factor and nephroblastoma-overexpressed protein. *PLoS ONE* 5: e16000.
- Aimo, L., et al. 2010. Gestational zinc deficiency affects the regulation of transcription factors AP-1, NF κ B and NFAT in fetal brain. *J. Nutr. Biochem.* 21: 1069-1075.
- Paulsson, A.K., et al. 2010. Post-translational regulation of calsarcin-1 during pressure overload-induced cardiac hypertrophy. *J. Mol. Cell. Cardiol.* 48: 1206-1214.
- Stühmer, T., et al. 2010. Preclinical anti-myeloma activity of the novel HDAC-inhibitor JNJ-26481585. *Br. J. Haematol.* 149: 529-536.
- Mallery, D.L., et al. 2010. Antibodies mediate intracellular immunity through tripartite motif-containing 21 (TRIM21). *Proc. Natl. Acad. Sci. USA* 107: 19985-19990.
- Alexandru, N., et al. 2011. Platelet activation in hypertension associated with hypercholesterolemia: effects of irbesartan. *J. Thromb. Haemost.* 9: 173-184.
- Arora, D., et al. 2011. Protein-tyrosine phosphatase DEP-1 controls receptor tyrosine kinase FLT3 signaling. *J. Biol. Chem.* 286: 10918-10929.