

ULBP3 (L-20): sc-20574

BACKGROUND

The immune system contains genetically encoded subsystems which monitor the extracellular environment in order to eliminate pathogens and resolve abnormal or transformed tissues. Cytomegalovirus UL16 binding proteins, known as ULBPs, are GPI-linked glycoproteins that belong to the extended MHC class I family and are distantly related to MHC class I polypeptide-related sequence B, known as MICB. ULBP and MICB proteins are ligands for the activating receptor, NKG2D/DAP10, which causes lymphocyte activation, resulting in the secretion of cytokines, such as interferon- γ , and tumor cell lysis. The interaction of ULBP or MICB with NKG2D/DAP10 can be blocked by the soluble form of UL16. ULBPs stimulate cytokine and chemokine production from NK cells, CD8 $\alpha\beta$ T cells and $\gamma\delta$ T cells. Soluble forms of ULBPs induce protein tyrosine phosphorylation and activation of the Janus kinase 2, Stat5, extracellular signal-regulated kinase, mitogen-activated protein kinase and phosphatidylinositol 3-kinase (PI 3-kinase)/Akt signal transduction pathways.

REFERENCES

1. Kubin, M., et al. 2001. ULBP1, 2, 3: novel MHC class I-related molecules that bind to human cytomegalovirus glycoprotein UL16, activate NK cells. *Eur. J. Immunol.* 31: 1428-1437.
2. Cosman, D., et al. 2001. ULBPs, novel MHC class I-related molecules, bind to CMV glycoprotein UL16 and stimulate NK cytotoxicity through the NKG2D receptor. *Immunity* 14: 123-133.
3. Steinle, A., et al. 2001. Interactions of human NKG2D with its ligands MICA, MICB and homologs of the mouse RAE-1 protein family. *Immunogenetics* 53: 279-287.
4. Sutherland, C.L., et al. 2002. UL16-binding proteins, novel MHC class I-related proteins, bind to NKG2D and activate multiple signaling pathways in primary NK cells. *J. Immunol.* 168: 671-679.
5. LocusLink Report (LocusID: 4277). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: ULBP3 (human) mapping to 6q25.1.

SOURCE

ULBP3 (L-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ULBP3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-20574 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ULBP3 (L-20) is recommended for detection of ULBP3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for ULBP3 siRNA (h): sc-43006, ULBP3 shRNA Plasmid (h): sc-43006-SH and ULBP3 shRNA (h) Lentiviral Particles: sc-43006-V.

Molecular Weight of ULBP3: 30 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ULBP3 (L-20): sc-20574. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **ULBP3 (D-1): sc-390844** or **ULBP3 (YY42): sc-80420**, our highly recommended monoclonal alternatives to ULBP3 (L-20).