

goat anti-mouse IgG_{2b}-HRP: sc-2062

BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. We offer horseradish peroxidase (HRP) and alkaline phosphatase (AP) conjugated isotype-specific secondary antibodies for detection of mouse IgG₁, IgG_{2a}, IgG_{2b}, IgG₃, IgM and IgA in Western Blot and ELISA. Pre-adsorbed HRP and AP conjugated isotype-specific secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples.

SOURCE

goat anti-mouse IgG_{2b}-HRP is an affinity purified secondary antibody raised in goat against mouse IgG_{2b} and conjugated to HRP (horseradish peroxidase).

PRODUCT

Each vial contains 200 µg IgG_{2b} in 0.5 ml of 1X PBS containing 40% glycerol.

APPLICATIONS

goat anti-mouse IgG_{2b}-HRP is recommended for detection of mouse IgG_{2b} by Western Blotting (starting dilution: 1:1000, dilution range 1:1,000-1:10,000; optimal dilution to be determined by titration) and immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400).


RECOMMENDED SUPPORT PRODUCTS

- UltraCruz™ Tissue Culture Dish, 100 mm polystyrene dish: sc-200286
- UltraCruz™ Cell Scrapers, 25 cm, sterile, 100 per case: sc-213229
- RIPA Lysis Buffer, 50 ml, cell lysis buffer with protease inhibitors: sc-24948
- Complete™ Protease Inhibitor Cocktail Tablet, 20 tablets: sc-291300
- Electrophoresis Sample Buffer, 2X, 25 ml, reducing buffer: sc-24945
- UltraCruz™ PVDF Transfer membrane, 0.45 µm, 30 cm x 3 m roll: sc-3723
- UltraCruz™ Nitrocellulose Pure Transfer Membrane, 0.22 µm, 30 cm x 3 m roll: sc-3718
- Cruz Blot-A: sc-3901 (Western Blotting membrane with human cell line extracts from 10 different cell types)
- Running Buffer, 10X, 1 L, TRIS-Glycine WB running buffer, pH 8.3: sc-24949
- Towbin, with SDS, 10X, 1 L, WB transfer buffer pH 8.3: sc-24954
- Bovine Serum Albumin (BSA), 100 g, blocking/incubation agent: sc-2323
- TBS Blotto A, lyophilized powder in single-use bottle: sc-2333
- Western Blotting Luminol Reagent, for 2,000 cm² membrane area: sc-2048
- UltraCruz™ Electrophoresis Cell: sc-201625 : runs up to 10 or 15 sample by SDS – PAGE protein electrophoresis
- UltraCruz™ Autoradiography Film, Blue, 8 x 1, 100 sheets: sc-201697
- CrystalCruz™ Micro Slides 75 x 25 mm; frosted sides: sc-24976
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



goat anti-mouse IgG_{2b}-HRP: sc-2062. Western blot analysis of ERK 2 expression in Jurkat whole cell lysate. Antibody tested: ERK 2 (D-2): sc-1647.

SELECT PRODUCT CITATIONS

1. Murphy, B.J., et al. 2005. The metal-responsive transcription factor-1 contributes to HIF-1 activation during hypoxic stress. *Biochem. Biophys. Res. Commun.* 337: 860-867.
2. Amin, K., et al. 2006. Characterization of ANIT-induced toxicity using precision-cut rat and dog liver slices cultured in a dynamic organ roller system. *Toxicol. Pathol.* 34: 776-784.
3. Page, T.J., et al. 2006. Genome-wide analysis of human HSF1 signaling reveals a transcriptional program linked to cellular adaptation and survival. *Mol. Biosyst.* 2: 627-639.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.