

# goat anti-mouse IgG<sub>2b</sub>-AP: sc-2068

## BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. We offer horseradish peroxidase (HRP) and alkaline phosphatase (AP) conjugated isotype-specific secondary antibodies for detection of mouse IgG<sub>1</sub>, IgG<sub>2a</sub>, IgG<sub>2b</sub>, IgG<sub>3</sub>, IgM and IgA in Western Blot and ELISA. Pre-adsorbed HRP and AP conjugated isotype-specific secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples.

## SOURCE

goat anti-mouse IgG<sub>2b</sub>-AP is an affinity purified secondary antibody raised in goat against mouse IgG<sub>2b</sub> and conjugated to AP (alkaline phosphatase).

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> in 0.5 ml of PBS containing 50% glycerol, 1 mM zinc chloride, 0.02% sodium azide and 1 mM magnesium chloride.

## APPLICATIONS

goat anti-mouse IgG<sub>2b</sub>-AP is recommended for detection of mouse IgG<sub>2b</sub> by Western Blotting (starting dilution: 1:1000, dilution range 1:1,000-1:10,000; optimal dilution to be determined by titration) and immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400).

## RECOMMENDED SUPPORT PRODUCTS

- UltraCruz™ Tissue Culture Dish, 100 mm polystyrene dish: sc-200286
- UltraCruz™ Cell Scrapers, 25 cm, sterile, 100 per case: sc-213229
- RIPA Lysis Buffer, 50 ml, cell lysis buffer with protease inhibitors: sc-24948
- Complete™ Protease Inhibitor Cocktail Tablet, 20 tablets: sc-291300
- Electrophoresis Sample Buffer, 2X, 25 ml, reducing buffer: sc-24945
- UltraCruz™ PVDF Transfer membrane, 0.45 µm, 30 cm x 3 m roll: sc-3723
- UltraCruz™ Nitrocellulose Pure Transfer Membrane, 0.22 µm, 30 cm x 3 m roll: sc-3718
- Cruz Blot-A: sc-3901 (Western Blotting membrane with human cell line extracts from 10 different cell types)
- Running Buffer, 10X, 1 L, TRIS-Glycine WB running buffer, pH 8.3: sc-24949
- Towbin, with SDS, 10X, 1 L, WB transfer buffer pH 8.3: sc-24954
- Bovine Serum Albumin (BSA), 100 g, blocking/incubation agent: sc-2323
- TBS Blotto A, lyophilized powder in single-use bottle: sc-2333
- Western Blotting Luminol Reagent, for 2,000 cm<sup>2</sup> membrane area: sc-2048
- UltraCruz™ Electrophoresis Cell: sc-201625 : runs up to 10 or 15 sample by SDS – PAGE protein electrophoresis
- UltraCruz™ Autoradiography Film, Blue, 8 x 1, 100 sheets: sc-201697
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973

## SELECT PRODUCT CITATIONS

1. Debeb, B.G., et al. 2009. Isolation of Oct4-expressing extraembryonic endoderm precursor cell lines. PLoS ONE 4: e7216.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.