

A cyclase II (H-100): sc-20761

BACKGROUND

Adenylyl cyclases function to convert ATP to cyclic AMP in response to activation by a variety of hormones, neurotransmitters and other regulatory molecules. Cyclic AMP, in turn, activates several other target molecules to control a broad range of diverse phenomena such as metabolism, gene transcription and memory. Adenylyl cyclases respond to receptor-initiated signals, mediated by the G_s and G_i heterotrimeric G proteins. The binding of an agonist to a G_s -coupled receptor catalyzes the exchange of GDP (bound to $G_{\alpha s}$) for GTP, the dissociation of GTP- $G_{\alpha s}$ from $G_{\beta\gamma}$ and $G_{\alpha s}$ -mediated activation of adenylyl cyclase. Adenylyl cyclases of the type II family differ from other subforms in that they are conditionally stimulated by $G_{\alpha s/\beta\gamma}$ subunits and regulated by PKC-mediated C-terminal phosphorylation. Both short- and long-term activation of D(2L) dopamine receptors result in a marked degree of sensitization of A cyclase I, II, V and IX, but not A cyclase VIII. The effects on A cyclase I, II and VIII is dependent upon the ability of these A cyclase isoforms to synergistically respond to selective activators in the presence of activated $G_{\alpha s}$.

REFERENCES

- Gilman, A.G. 1987. G proteins: transducers of receptor-generated signals. *Ann. Rev. Biochem.* 56: 615-649.
- Bourne, H.R., et al. 1990. The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348: 125-132.
- Tang, W.J., et al. 1992. Adenylyl cyclases. *Cell* 70: 869-872.
- Taussig, R., et al. 1994. Distinct patterns of bidirectional regulation of mammalian adenylyl cyclases. *J. Biol. Chem.* 269: 6093-6100.
- Jacobowitz, O., et al. 1994. Phorbol ester-induced stimulation and phosphorylation of adenylyl cyclase 2. *Proc. Natl. Acad. Sci. USA* 91: 10630-10634.
- Bol, G.F., et al. 1997. Adenylyl cyclase type II is stimulated by PKC via C-terminal phosphorylation. *Biochim. Biophys. Acta* 1358: 307-313.
- Cumbay, M.G., et al. 2001. Heterologous sensitization of recombinant adenylyl cyclases by activation of D2 dopamine receptors. *J. Pharmacol. Exp. Ther.* 297: 1201-1209.

CHROMOSOMAL LOCATION

Genetic locus: ADCY2 (human) mapping to 5p15.31; Adcy2 (mouse) mapping to 13 B3.

SOURCE

A cyclase II (H-100) is a rabbit polyclonal antibody raised against amino acids 441-540 of A cyclase II of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

A cyclase II (H-100) is recommended for detection of A cyclase II of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

A cyclase II (H-100) is also recommended for detection of A cyclase II in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for A cyclase II siRNA (h): sc-40317, A cyclase II siRNA (m): sc-40318, A cyclase II shRNA Plasmid (h): sc-40317-SH, A cyclase II shRNA Plasmid (m): sc-40318-SH, A cyclase II shRNA (h) Lentiviral Particles: sc-40317-V and A cyclase II shRNA (m) Lentiviral Particles: sc-40318-V.

Molecular Weight of A cyclase II: 124 kDa.

Positive Controls: SH-SY5Y cell lysate: sc-3812 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

- Bek, M.J., et al. 2001. Differential expression of adenylyl cyclases in the rat nephron. *Kidney Int.* 60: 890-899.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **A cyclase II (F-7): sc-514938**, our highly recommended monoclonal alternative to A cyclase II (H-100).