# TNP2 (R-19): sc-21129



The Power to Question

#### **BACKGROUND**

During mammalian spermiogenesis, histones are transiently replaced by several low molecular weight proteins called transition proteins (TNPs). Transition proteins facilitate chromatin transformation from the nucleosome structure to the nucleoprotamine structure during spermatid differentiation. Transition protein-2, also known as TNP2, and TP2, maps to human chromosome 16p13.13 and encodes a highly basic nuclear protein. TNP2 is a spermatid-specific product of the haploid genome which replaces histone and is itself replaced in the mature sperm by the protamines. TNP2 is not a critical factor for shaping of the sperm nucleus, histone displacement, initiation of chromatin condensation, binding of protamines to DNA, or fertility. However, TNP2 is necessary for maintaining the normal processing of protamine 2 and, consequently, the completion of chromatin condensation. If TNP1 is missing, TNP2 may partially compensate for TNP1, but this dysregulation of nucleoprotein replacement results in an abnormal pattern of chromatin condensation and in reduced fertility.

## REFERENCES

- Nelson, J., et al. 1993. Linkage of human spermatid-specific basic nuclear protein genes. Definition and evolution of the P1-->P2-->TP2 locus. J. Biol. Chem. 268: 2932-2936.
- 2. Online Mendelian Inheritance in Man, OMIM™. Johns Hopkins University, Baltimore, MD. MIM Number: 190232: 7/13/1999: World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- 3. Yu, Y., et al. 2000. Anormal spermatogenisis and reduced fertility in transition nuclear protein 1-deficient mice. Proc. Nat. Acad. Sci. USA 97: 4683-4688.
- 4. Online Mendelian Inheritance in Man, OMIM™. Johns Hopkins University, Baltimore, MD. MIM Number: 190231: 7/19/2000: World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Zhao, M., et al. 2001. Targeted disruption of the transition protein 2 gene affects sperm chromatin structure and reduces fertility in mice. Mol. Cell. Biol. 21: 7243-7255.
- 6. LocusLink Report (LocusID: 7142). http://www.ncbi.nlm.nih.gov/LocusLink/

## **SOURCE**

TNP2 (R-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TNP2 of rat origin.

#### **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-21129 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **APPLICATIONS**

TNP2 (R-19) is recommended for detection of TNP2 of rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2  $\mu$ g per 100–500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of TNP2: 13 kDa.

Positive Controls: rat testes extract: sc-2400.

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/ 2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

# DATA



TNP2 (R-19): sc-21129. Western blot analysis of TNP2 expression in rat testis tissue extract.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

## **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3801 **fax** 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.co**