

V-ATPase B1 (N-20): sc-21206

BACKGROUND

Vacuolar-type H⁺-ATPase (V-ATPase) is a multisubunit enzyme responsible for acidification of eukaryotic intracellular organelles. V-ATPases pump protons against an electrochemical gradient, while F-ATPases reverse the process, thereby synthesizing ATP. A peripheral V₁ domain, which is responsible for ATP hydrolysis, and an integral V₀ domain, which is responsible for proton translocation, compose V-ATPase. Nine subunits (A-H) make up the V₁ domain and five subunits (a, d, c, c' and c'') make up the V₀ domain. Like F-ATPase, V-ATPase most likely operates through a rotary mechanism. The V-ATPase V₁ B subunit exists as two isoforms. In the inner ear, the V-ATPase B1 isoform functions in proton secretion and is required to maintain proper endolymph pH and normal auditory function. The gene encoding the human V-ATPase B1 isoform maps to chromosome 2q13.3. Mutations in this gene cause distal renal tubular acidosis associated with sensorineural deafness. The V-ATPase B2 isoform is expressed in kidney and is the only B isoform expressed in osteoclasts. The gene encoding the human V-ATPase B2 isoform maps to chromosome 8p22-p21.

REFERENCES

- Bernasconi, P., et al. 1990. An mRNA from human brain encodes an isoform of the B subunit of the vacuolar H⁺-ATPase. *J. Biol. Chem.* 265: 17428-17431.
- Ozcelik, T., et al. 1991. Chromosomal assignments of genes for vacuolar (endomembrane) proton pump subunits VPP1/Vpp-1 (116 kDa) and VPP3/Vpp-3 (58 kDa) in human and mouse. *Cytogenet. Cell Genet.* 58: 2008-2009.
- Nelson, R.D., et al. 1992. Selectively amplified expression of an isoform of the vacuolar H⁺-ATPase 56 kilodalton subunit in renal intercalated cells. *Proc. Natl. Acad. Sci. USA* 89: 3541-3545.
- Lee, B.S., et al. 1996. Osteoclasts express the B2 isoform of vacuolar H⁺-ATPase intracellularly and on their plasma membranes. *Am. J. Physiol.* 270: 382-388.
- Karet, F.E., et al. 1999. Mutations in the gene encoding B1 subunit of H⁺-ATPase cause renal tubular acidosis with sensorineural deafness. *Nat. Genet.* 21: 84-90.
- Nishi, T., et al. 2002. The vacuolar H⁺-ATPases—nature's most versatile proton pumps. *Nat. Rev. Mol. Cell. Biol.* 3: 94-103.

CHROMOSOMAL LOCATION

Genetic locus: ATP6V1B1 (human) mapping to 2p13.3; Atp6v1b1 (mouse) mapping to 6 C3.

SOURCE

V-ATPase B1 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of V-ATPase B1 of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-21206 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

V-ATPase B1 (N-20) is recommended for detection of V-ATPase B1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


V-ATPase B1 (N-20) is also recommended for detection of V-ATPase B1 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for V-ATPase B1 siRNA (h): sc-36787, V-ATPase B1 siRNA (m): sc-36788, V-ATPase B1 shRNA Plasmid (h): sc-36787-SH, V-ATPase B1 shRNA Plasmid (m): sc-36788-SH, V-ATPase B1 shRNA (h) Lentiviral Particles: sc-36787-V and V-ATPase B1 shRNA (m) Lentiviral Particles: sc-36788-V.


Molecular Weight of V-ATPase B1: 56-58 kDa.

Positive Controls: mouse kidney extract: sc-2255 or rat kidney extract: sc-2394.

DATA



V-ATPase B1 (N-20): sc-21206. Western blot analysis of V-ATPase B1 expression in mouse kidney (A) and rat kidney (B) tissue extracts.



V-ATPase B1 (N-20): sc-21206. Immunoperoxidase staining of formalin fixed, paraffin-embedded human kidney tissue showing cytoplasmic staining of cells in tubules.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **V-ATPase B1/2 (D-4): sc-271832** or **V-ATPase B1/2 (F-6): sc-55544**, our highly recommended monoclonal alternatives to V-ATPase B1 (N-20).