

V-ATPase H (N-20): sc-21228

BACKGROUND

Vacuolar-type H⁺-ATPase (V-ATPase) is a multisubunit enzyme responsible for acidification of eukaryotic intracellular organelles. V-ATPases pump protons against an electrochemical gradient, while F-ATPases reverse the process, thereby synthesizing ATP. A peripheral V₁ domain, which is responsible for ATP hydrolysis, and an integral V₀ domain, which is responsible for proton translocation, compose V-ATPase. Nine subunits (A-H) make up the V₁ domain and five subunits (a, d, c, c' and c'') make up the V₀ domain. Like F-ATPase, V-ATPase most likely operates through a rotary mechanism. The H subunit of V-ATPase, also designated SDF is comprised of two polypeptides derived from the same gene. This regulatory subunit plays a critical role in the functional coupling of ATP hydrolysis activity to proton transport in the V-ATPase pump.

CHROMOSOMAL LOCATION

Genetic locus: ATP6V1H (human) mapping to 8q11.23; Atp6v1h (mouse) mapping to 1 A1.

SOURCE

V-ATPase H (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of V-ATPase H of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-21228 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

V-ATPase H (N-20) is recommended for detection of V-ATPase H of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

V-ATPase H (N-20) is also recommended for detection of V-ATPase H in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for V-ATPase H siRNA (h): sc-36801, V-ATPase H siRNA (m): sc-36802, V-ATPase H shRNA Plasmid (h): sc-36801-SH, V-ATPase H shRNA Plasmid (m): sc-36802-SH, V-ATPase H shRNA (h) Lentiviral Particles: sc-36801-V and V-ATPase H shRNA (m) Lentiviral Particles: sc-36802-V.


Molecular Weight of V-ATPase H: 50/57 kDa.

Positive Controls: Caki-1 cell lysate: sc-2224, SK-N-SH cell lysate: sc-2410 or V-ATPase H (m): 293T Lysate: sc-124526.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



V-ATPase H (N-20): sc-21228. Western blot analysis of V-ATPase H expression in non-transfected: sc-117752 (A) and mouse V-ATPase H transfected: sc-124526 (B) 293T whole cell lysates.



V-ATPase H (N-20): sc-21228. Immunofluorescence staining of methanol-fixed Hep G2 cells showing cytoplasmic localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **V-ATPase H (C-8): sc-166227** or **V-ATPase H (E-10): sc-271186**, our highly recommended monoclonal alternatives to V-ATPase H (N-20).