

FOXC1 (N-19): sc-21394

BACKGROUND

The forkhead transcription factor genes FOXC1 (Mf1) and FOXC2 (Mfh1) interact with the Notch signaling pathway and are required for the pre patterning of anterior and posterior domains in the presumptive somites through a putative Notch/ Δ /Mesp regulatory loop. The genes have similar, dose-dependent functions, and compensate for each other in the early development of the heart, blood vessels and somites. Both FOXC1 and FOXC2 are expressed in the mesenchyme from which the ocular drainage structures derive. FOXC1 and FOXC2 also interact in kidney and heart development. Mutations in the FOXC1 gene result in Axenfeld-Rieger malformations of the anterior segment of the eye and lead to an increased susceptibility of glaucoma, including juvenile glaucoma. Functional regions in FOXC1 are required for nuclear localization and transcriptional regulation. Specifically, two regions in the FOXC1 forkhead domain, one rich in basic amino acid residues, and a second, highly conserved among all FOX proteins, are necessary for nuclear localization of the FOXC1 protein.

REFERENCES

- Smith, R.S., Zabaleta, A., Kume, T., Savinova, O.V., Kidson, S.H., Martin, J.E., Nishimura, D.Y., Alward, W.L., Hogan, B.L. and John, S.W. 2000. Haploinsufficiency of the transcription factors FOXC1 and FOXC2 results in aberrant ocular development. *Hum. Mol. Genet.* 9: 1021-1032.
- Kume, T., Deng, K. and Hogan, B.L. 2000. Murine forkhead/winged helix genes FOXC1 (Mf1) and FOXC2 (Mfh1) are required for the early organogenesis of the kidney and urinary tract. *Development* 127: 1387-1395.
- Kawase, C., Kawase, K., Taniguchi, T., Sugiyama, K., Yamamoto, T., Kitazawa, Y., Alward, W.L., Stone, E.M., Nishimura, D.Y. and Sheffield, V.C. 2001. Screening for mutations of Axenfeld-Rieger syndrome caused by FOXC1 gene in Japanese patients. *J. Glaucoma* 10: 477-482.
- Kume, T., Jiang, H., Topczewska, J.M. and Hogan, B.L. 2001. The murine winged helix transcription factors, FOXC1 and FOXC2, are both required for cardiovascular development and somitogenesis. *Genes Dev.* 15: 2470-2482.
- Berry, F.B., Saleem, R.A. and Walter, M.A. 2002. FOXC1 transcriptional regulation is mediated by N- and C-terminal activation domains and contains a phosphorylated transcriptional inhibitory domain. *J. Biol. Chem.* 277: 10292-10297.

CHROMOSOMAL LOCATION

Genetic locus: FOXC1 (human) mapping to 6p25.3; Foxc1 (mouse) mapping to 13 A3.2.

SOURCE

FOXC1 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of FOXC1 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-21394 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

FOXC1 (N-19) is recommended for detection of FOXC1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for FOXC1 siRNA (h): sc-43766, FOXC1 siRNA (m): sc-145221, FOXC1 shRNA Plasmid (h): sc-43766-SH, FOXC1 shRNA Plasmid (m): sc-145221-SH, FOXC1 shRNA (h) Lentiviral Particles: sc-43766-V and FOXC1 shRNA (m) Lentiviral Particles: sc-145221-V

Positive Controls: K-562 whole cell lysate: sc-2203.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



FOXC1 (N-19): sc-21394. Western blot analysis of FOXC1 expression in K-562 whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **FOXC1 (4D11): sc-293455**, our highly recommended monoclonal alternative to FOXC1 (N-19).