

Glut10 (L-20): sc-21635

BACKGROUND

Glucose serves as the major energy substrate of mammalian cells and is fundamental to metabolism. Glucose passage across cell membranes is mediated by a family of transporters termed glucose transporters, or Gluts, which are characterized by the presence of 12 membrane-spanning helices. The Glut family is divided into three subfamilies: class I (previously known as glucose transporters), which includes Glut1, Glut2, Glut3 and Glut4; class II (previously known as fructose transporters), which includes Glut5, Glut7, Glut9 and Glut11; and class III, which includes Glut6, Glut8, Glut10, Glut12 and the myo-inositol transporter HMIT1. Glut10 is a 541 amino acid facilitative glucose transporter that is highly expressed in liver and pancreas. It contains 12 transmembrane domains, with a hydrophilic intracellular loop between helices 6 and 7, and a potential N-linked glycosylation site, with a large extracellular loop between helices 9 and 10. The gene for Glut 10, SLC2A10, maps to chromosome 20q13.12, a region that is linked to type 2 diabetes.

CHROMOSOMAL LOCATION

Genetic locus: SLC2A10 (human) mapping to 20q13.12; Slc2a10 (mouse) mapping to 2 H3.

SOURCE

Glut10 (L-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of Glut10 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-21635 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Glut10 (L-20) is recommended for detection of Glut10 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Glut10 siRNA (h): sc-60697, Glut10 siRNA (m): sc-60698, Glut10 shRNA Plasmid (h): sc-60697-SH, Glut10 shRNA Plasmid (m): sc-60698-SH, Glut10 shRNA (h) Lentiviral Particles: sc-60697-V and Glut10 shRNA (m) Lentiviral Particles: sc-60698-V.


Molecular Weight of Glut10: 57 kDa.

Positive Controls: MDA-MB-435S whole cell lysate: sc-364184, HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



Glut10 (L-20): sc-21635. Western blot analysis of Glut10 expression in MDA-MB-435S (A) and HeLa (B) whole cell lysates.



Glut10 (L-20): sc-21635. Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing cytoplasmic staining of exocrine glandular cells.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Glut10 (H-10): sc-398495**, our highly recommended monoclonal alternative to Glut10 (L-20).