SANTA CRUZ BIOTECHNOLOGY, INC.

c-GMP (8D8): sc-21727



BACKGROUND

The Golgi complex plays an essential role in the post-translational modification and sorting of proteins transported from the endoplasmic reticulum (ER). The Golgi stack consists of a distinct *cis* face, or entry face, and a *trans* face, or exit face, which are connected via the *cis*, medial and *trans* Golgi networks. The networks are functionally distinct; different enzymes are contained within each compartment and impart different posttranslational modifications (glycosylation, sulfation, phosphorylation, acylation and proteolytic digestion) on proteins as they are transported through the organelle. The 130 kDa c-GMP *(cis*-Golgi membrane protein) localizes to both the *cis* and medial cisternae, whereas t-GMP (100 kDa) localizes to the *trans*-most cisternae and the *trans*tubular network. Treatment with the fungal metabolite brefeldin A (BFA) induces both disassembly of the *cis/*middle- and *trans*-Golgi complex and also translocation of c-GMP and t-GMP to the ER. In the adult rat epididymis, t-GMP and c-GMP exhibit a reticular, perinuclear pattern *in vitro* expression pattern.

REFERENCES

- Barriocanal, J.G., et al. 1986. Biosynthesis, glycosylation, movement through the Golgi system, and transport to lysosomes by an N-linked carbohydrate-independent mechanism of three lysosomal integral membrane proteins. J. Biol. Chem. 261: 16755-16763.
- Yuan, L., et al. 1987. Two integral membrane proteins located in the *cis*middle and *trans*-part of the Golgi system acquire sialylated N-linked carbohydrates and display different turnovers and sensitivity to cAMPdependent phosphorylation. J. Cell Biol. 105: 215-227.
- Alcalde, J., et al. 1992. Assembly and disassembly of the Golgi complex: two processes arranged in a *cis-trans* direction. J. Cell Biol. 116: 69-83.
- Suarez-Quian, C.A. and Jelesoff, N. 1994. Two Golgi integral membrane proteins (GIMPS) exhibit region- and cell type-specific distribution in the epididymis of the adult rat. Microsc. Res. Tech. 29: 481-491.

SOURCE

c-GMP (8D8) is a mouse monoclonal antibody raised against integral membrane protein from liver fraction enriched in Golgi cisternae of rat origin.

PRODUCT

Each vial contains 200 μg lgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

c-GMP (8D8) is available conjugated to agarose (sc-21727 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-21727 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-21727 PE), fluorescein (sc-21727 FITC), Alexa Fluor[®] 488 (sc-21727 AF488), Alexa Fluor[®] 546 (sc-21727 AF546), Alexa Fluor[®] 594 (sc-21727 AF594) or Alexa Fluor[®] 647 (sc-21727 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-21727 AF680) or Alexa Fluor[®] 790 (sc-21727 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

c-GMP (8D8) is recommended for detection of c-GMP of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of c-GMP: 130 kDa.

Positive Controls: KNRK whole cell lysate: sc-2214 or 3611-RF whole cell lysate: sc-2215.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



c-GMP (8D8): sc-21727. Western blot analysis of c-GMP expression in KNRK (A) and 3611-RF (B) cell extracts.

SELECT PRODUCT CITATIONS

- Jurdana, M., et al. 2009. Neural Agrin changes the electrical properties of developing human skeletal muscle cells. Cell. Mol. Neurobiol. 29: 123-131.
- Zhang, L., et al. 2021. Cardioprotective effect of icariin against myocardial fibrosis and its molecular mechanism in diabetic cardiomyopathy based on network pharmacology: role of ICA in DCM. Phytomedicine 91: 153607.
- Jin, M., et al. 2022. Tianma Gouteng decoction exerts pregnancy-protective effects against preeclampsia via regulation of oxidative stress and NO signaling. Front. Pharmacol. 13: 849074.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.