SANTA CRUZ BIOTECHNOLOGY, INC.

CD53 (202-24b): sc-21778



BACKGROUND

The CD53 antigen is a member of the tetraspanin membrane protein family that is expressed in the lymphoid-myeloid lineage. The tetraspanin superfamily (CD9, CD37, CD53, CD63, CD81 and CD82) comprises a group of cell-surface proteins that are involved in cell activation and signal transduction as well as in cell adhesion, motility and metastasis. Tetraspanin transmembrane proteins have a metastasis suppressor effect by acting as cell motility brakes in tumor cells. Human neutrophils express high levels of CD53, an N-glycosylated pan-leukocyte antigen and the true homolog of the rat MRC-OX-44 antigen. CD53 is expressed on B cells, monocytes, macrophages, neutrophils, single (CD4 or CD8) positive thymocytes and peripheral T cells.

REFERENCES

- Bazil, V., et al. 1989. Monoclonal antibodies against human leucocyte antigens. III. Antibodies against CD45R, CD6, CD44 and two newly described broadly expressed glycoproteins MEM-53 and MEM-102. Folia Biol. 35: 289-297.
- Angelisova, P., et al. 1990. The human leucocyte surface antigen CD53 is a protein structurally similar to the CD37 and MRC OX-44 antigens. Immunogenetics 32: 281-285.
- Olweus, J., et al. 1993. CD53, a protein with four membrane-spanning domains, mediates signal transduction in human monocytes and B cells. J. Immunol. 151: 707-716.
- Rasmussen, A.M., et al. 1994. Cross-linking of CD53 promotes activation of resting human B lymphocytes. J. Immunol. 153: 4997-5007.
- 5. Okochi, H., et al. 1997. Expression of tetraspans transmembrane family (CD9, CD37, CD53, CD63, CD81 and CD82) in normal and neoplastic human keratinocytes: an association of CD9 with α 3 β 1 Integrin. Br. J. Dermatol. 137: 856-863.
- Lagaudrière-Gesbert, C., et al. 1997. Functional analysis of four tetraspans, CD9, CD53, CD81, and CD82, suggests a common role in costimulation, cell adhesion, and migration: only CD9 upregulates HB-EGF activity. Cell. Immunol. 182: 105-112.
- Mollinedo, F., et al. 1998. Physiological activation of human neutrophils down-regulates CD53 cell surface antigen. J. Leukoc. Biol. 63: 699-706.
- Beinert, T., et al. 2000. Increased expression of the tetraspanins CD53 and CD63 on apoptotic human neutrophils. J. Leukoc. Biol. 67: 369-373.
- Yunta, M., et al. 2002. Transient activation of the c-Jun N-terminal kinase (JNK) activity by ligation of the tetraspan CD53 antigen in different cell types. Eur. J. Biochem. 269: 1012-1021.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: CD53 (human) mapping to 1p13.3.

SOURCE

CD53 (161-2) is a mouse monoclonal antibody raised against stimulated human leukocytes.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CD53 (202-24b) is available conjugated to either phycoerythrin (sc-21778 PE) or fluorescein (sc-21778 FITC), 200 μ g/ml, for IF, IHC(P) and FCM.

APPLICATIONS

CD53 (202-24b) is recommended for detection of CD53 of human origin by flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for CD53 siRNA (h): sc-42796, CD53 shRNA Plasmid (h): sc-42796-SH and CD53 shRNA (h) Lentiviral Particles: sc-42796-V.

Molecular Weight of CD53: 32-45 kDa.

DATA





CD53 (202-24b) PE: sc-21778 PE. FCM analysis of human peripheral blood leukocytes. Black line histogram represents the isotype control, normal mouse IgG₁-PE: sc-2866. CD53 (202-24b) FITC: sc-21778 FITC. FCM analysis of human peripheral blood leukocytes. Black line histogram represents the isotype control, normal mouse lgG_1 -FITC: sc-2855.

RESEARCH USE

For research use only, not for use in diagnostic procedures.