

# Tropomyosin (D-17): sc-21806

## BACKGROUND

Tropomyosins are a group of structural proteins. Tropomyosins are present in virtually all eukaryotic cells (both muscle and nonmuscle), where they bind actin filaments and function to modulate actin-myosin interaction and stabilize actin filament structure. Tropomyosin  $\alpha$  is encoded by the TPM1 gene, which maps to human chromosome 15q22.2 and undergoes alternative splicing to generate at least four isoforms including skeletal muscle (isoform 1), smooth muscle (isoform 2), fibroblast/TM3 (isoform 3) and isoform 4. Tropomyosin  $\beta$  is encoded by the TPM2 gene, which maps to human chromosome 9p13.3 and undergoes alternative splicing to generate three isoforms including skeletal muscle (isoform 1), nonmuscle/fibroblast TM36/epithelial TMe1 (isoform 2) and nonmuscle (isoform 3). Troponin I binds Tropomyosin at a specific region and the association of Tropomyosin-Troponin with actin filaments may increase the rigidity of actin filaments. Tropomyosin also interacts with caldesmon to regulate smooth muscle contraction.

## SOURCE

Tropomyosin (D-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Tropomyosin  $\beta$  of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-21806 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

Tropomyosin (D-17) is recommended for detection of Tropomyosin  $\alpha$  isoforms 1-4, Tropomyosin  $\beta$  isoforms 1-3 and Tropomyosin  $\gamma$  of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Tropomyosin (D-17) is also recommended for detection of Tropomyosin  $\alpha$  isoforms 1-4, Tropomyosin  $\beta$  isoforms 1-3 and Tropomyosin  $\gamma$  in additional species, including equine, canine, bovine, porcine and avian.

Molecular Weight (predicted) of Tropomyosin  $\alpha$ : 33 kDa.

Molecular Weight (predicted) of Tropomyosin  $\beta$ : 33 kDa.

Molecular Weight (predicted) of Tropomyosin  $\gamma$ : 33 kDa.

Molecular Weight (predicted) of Tropomyosin 4: 29 kDa.


Molecular Weight (observed) of Tropomyosin: 31-47 kDa.

Positive Controls: Sol8 cell lysate: sc-2249, L8 cell lysate: sc-3807 or CCD-1064Sk cell lysate: sc-2263.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



Tropomyosin (D-17): sc-21806. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic and intercalated disc staining of myocytes.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **Tropomyosin (F-6): sc-74480** or **Tropomyosin (C-3): sc-376339**, our highly recommended monoclonal alternatives to Tropomyosin (D-17).