

Cdk7 (dV-15): sc-22119

BACKGROUND

Progression through the cell cycle requires activation of a series of enzymes designated cyclin dependent kinases (Cdks). The monomeric catalytic subunit Cdk2, a critical enzyme for initiation of cell cycle progression, is completely inactive. Partial activation is achieved by the binding of regulatory cyclins such as cyclin D1, while full activation requires additional phosphorylation at Thr 160. The enzyme responsible for the phosphorylation of Cdk2 on Thr 160 and also of Cdc2 p34 on Thr 161, designated Cdk-activating kinase (CAK), has been partially purified and shown to be comprised of a catalytic subunit and a regulatory subunit. The catalytic subunit, designated Cdk7, has been identified as the mammalian homolog of MO15, a protein kinase demonstrated in starfish and *Xenopus*. The regulatory subunit is a novel cyclin (cyclin H) and is required for activation of Cdk7. Like other Cdks, Cdk7 contains a conserved threonine residue required for full activity; mutation of this residue severely reduces CAK activity.

REFERENCES

1. Nurse, P. 1994. Ordering S phase and M phase in the cell cycle. *Cell* 79: 547-550.
2. Sherr, C.J. 1994. G₁ phase progression: cycling on cue. *Cell* 79: 551-555.
3. Helchman, K.A., et al. 1994. Rules to replicate by. *Cell* 79: 557-562.
4. King, R.W., et al. 1994. Mitosis in transition. *Cell* 79: 563-571.
5. Hunter, T., et al. 1994. Cyclins and cancer II: cyclin D and Cdk inhibitors come of age. *Cell* 79: 573-582.
6. Kato, J.Y., et al. 1994. Regulation of cyclin D-dependent kinase 4 (Cdk4) by Cdk4-activating kinase. *Mol. Cell. Biol.* 14: 2713-2721.
7. Matsuoka, M., et al. 1994. Activation of cyclin-dependent kinase 4 (Cdk4) by mouse MO15-associated kinase. *Mol. Cell. Biol.* 14: 7265-7275.
8. Levedakou, E.N., et al. 1994. Two novel human serine/threonine kinases with homologies to the cell cycle regulating *Xenopus* MO15, and NIMA kinases: cloning and characterization of their expression pattern. *Oncogene* 9: 1977-1988.
9. Wu, L., et al. 1994. Molecular cloning of the human CAK1 gene encoding a cyclin-dependent kinase-activating kinase. *Oncogene* 9: 2089-2096.

SOURCE

Cdk7 (dV-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Cdk7 of *Drosophila melanogaster* origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-22119 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Cdk7 (dV-15) is recommended for detection of Cdk7 of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of Cdk7 isoforms: 42/37 kDa.

Positive Controls: Schneider's *Drosophila* line 2 whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Cdk7 (dV-15): sc-22119. Western blot analysis of Cdk7 expression in Schneider's *Drosophila* line 2 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.