

Na⁺ CP type III α (S-15): sc-22202

BACKGROUND

Voltage-gated sodium channels are selective ion channels that regulate the permeability of sodium ions in excitable cells. During the propagation of an action potential, sodium channels allow an influx of sodium ions, which rapidly depolarize the cell. The three glycoproteins that comprise the voltage-gated sodium channel proteins include a pore-forming α subunit, a non-covalently associated β 1 subunit and a disulfide-linked β 2 subunit. The two β subunits regulate the level of channel expression, modulate gating and function as cell adhesion molecules for cellular aggregation and cytoskeleton interaction. The α subunits of sodium channels type I and III are predominantly expressed in neuronal cell bodies and proximal processes, while type II α subunits are more abundant along axons. The β 1 subunit of sodium channel type I is expressed in brain, skeletal and cardiac muscle. In the brain, β 1 and β 2 are highly expressed in Purkinje cells, and β 1 is also expressed in the pyramidal cells of the deep cerebellar nuclei. Impaired voltage-gated sodium channels lead to a number of diseases including myotonia.

CHROMOSOMAL LOCATION

Genetic locus: SCN3A (human) mapping to 2q24.3; Scn3a (mouse) mapping to 2 C1.3.

SOURCE

Na⁺ CP type III α (S-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Na⁺ CP type III α of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-22202 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Na⁺ CP type III α (S-15) is recommended for detection of sodium channel type III α of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Na⁺ CP type III α (S-15) is also recommended for detection of sodium channel type III α in additional species, including equine, canine, bovine, porcine and avian.


Suitable for use as control antibody for Na⁺ CP type III α siRNA (h): sc-43955, Na⁺ CP type III α siRNA (m): sc-42647, Na⁺ CP type III α shRNA Plasmid (h): sc-43955-SH, Na⁺ CP type III α shRNA Plasmid (m): sc-42647-SH, Na⁺ CP type III α shRNA (h) Lentiviral Particles: sc-43955-V and Na⁺ CP type III α shRNA (m) Lentiviral Particles: sc-42647-V.

Molecular Weight of Na⁺ CP type III α : 227 kDa.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz[™]: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



Na⁺ CP type III α (S-15): sc-22202. Western blot analysis of Na⁺ CP type III α expression in EOC 20 whole cell lysate (A) and mouse embryo tissue extract (B).



Na⁺ CP type III α (S-15): sc-22202. Immunoperoxidase staining of formalin fixed, paraffin-embedded human gall bladder tissue showing membrane and cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

- Pinto, F.M., et al. 2009. Molecular and functional characterization of voltage-gated sodium channels in human sperm. *Reprod. Biol. Endocrinol.* 7: 71.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Na⁺ CP type III α (3F3): sc-517010**, our highly recommended monoclonal alternative to Na⁺ CP type III α (S-15).