

β-1,4-Gal-T3 (C-17): sc-22285

BACKGROUND

Several oligosaccharide structures and protein glycoconjugate types are found in nature. Homologous glycosyltransferase (GT) gene families catalyze the formation of glycosidic linkages. The β-1,3 galactosyltransferase (β3GalT) gene family encodes a set of type II transmembrane glycoproteins that are catalytically diverse and use different donor substrates (UDP-galactose and UDP-N-acetylglucosamine) and different acceptor sugars (N-acetylglucosamine, galactose, N-acetylgalactosamine) to catalyze the addition of an activated monosaccharide to a terminal lactose. The protein coding sequences for β-1,3-Gal-T genes comprise a single exon and are distantly related to the *Drosophila Brainiac* gene. The β-1,4-galactosyltransferase (β4GalT) gene family encodes type II membrane-bound glycoproteins that show exclusive specificity for the donor substrate, UDP-galactose. β-1,4Gal-T genes transfer galactose in a β-1,4 linkage to similar acceptor sugars; each gene has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. GTs on the surface of sperm in part mediate gamete adhesion by binding to appropriate carbohydrate substrates in the egg zona pellucida. In several tissues and cell lines, GTs localize to the Golgi complex.

REFERENCES

- Shur, B.D. 1984. The receptor function of galactosyltransferase during cellular interactions. *Mol. Cell. Biochem.* 61: 143-158.
- Shur, B.D. 1986. The receptor function of galactosyltransferase during mammalian fertilization. *Adv. Exp. Med. Biol.* 207: 79-93.
- Strous, G.J. 1986. Golgi and secreted galactosyltransferase. *CRC Crit. Rev. Biochem.* 21: 119-151.
- Amado, M., et al. 1998. A family of human beta3-galactosyltransferases. Characterization of four members of a UDP-galactose:beta-N-acetylglucosamine/beta-nacetyl-galactosamine beta-1,3-galactosyltransferase family. *J. Biol. Chem.* 273: 12770-12778.
- Amado, M., et al. 1999. Identification and characterization of large galactosyltransferase gene families: galactosyltransferases for all functions. *Biochim. Biophys. Acta* 1473: 35-53.
- Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: 603093: 05/16/2001. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- LocusLink Report (LocusID: 8708). <http://www.ncbi.nlm.nih.gov/LocusLink/>
- LocusLink Report (LocusID: 2683). <http://www.ncbi.nlm.nih.gov/LocusLink/>

SOURCE

β-1,4-Gal-T3 (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of β-1,4-galactosyltransferase 3 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-22285 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

β-1,4-Gal-T3 (C-17) is recommended for detection of β-1,4-galactosyltransferase 3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2 µg per 100–500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of β-1,4-Gal-T3: 44 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, HeLa whole cell lysate: sc-2200 or COLO 320DM cell lysate: sc-2226.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/ 2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



β-1,4-Gal-T3 (C-17): sc-22285. Western blot analysis of β-1,4-Gal-T3 expression in HL-60 (A), HeLa (B), COLO 320DM (C) and JEG-3 (D) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.