

# KIR7.1 (N-16): sc-22438

## BACKGROUND

G protein-coupled inwardly rectifying potassium channels (KIR3.1 through KIR3.4) are coupled to numerous neurotransmitter receptors in the brain and are abundantly expressed in the olfactory bulb, hippocampus, neocortex, dentate gyrus, cerebellar cortex and thalamus regions of the brain. Also known as GIRK, KIR3 potassium channels localize to the soma and dendrites as well as axons of neurons. Liberated  $G_{\beta\gamma}$  subunits from G protein heterotrimers bind to and regulate KIR3 channel activity.  $G_{\beta 3^-}$  and  $G_{\beta 4^-}$ -containing  $G_{\beta\gamma}$  dimers bind directly to cytoplasmic domains of KIR3 proteins and increase the  $K^+$  current while  $G_{\beta 5^-}$ -containing  $G_{\beta\gamma}$  dimers inhibit KIR3  $K^+$  current. KIR3 activity is also inhibited by tyrosine phosphorylation. Brain-derived neurotrophic factor activates receptor tyrosine kinase B, which then phosphorylates KIR3 tyrosine residues, effectively inactivating the KIR3 channels.

## REFERENCES

- Braun, A.P., et al. 1992. Activation of  $\alpha$  1-adrenoceptors modulates the inwardly rectifying potassium currents of mammalian atrial myocytes. *Pflugers Arch.* 421: 431-439.
- Ponce, A., et al. 1996. G protein-gated inward rectifier  $K^+$  channel proteins (GIRK1) are present in the soma and dendrites as well as in nerve terminals of specific neurons in the brain. *J. Neurosci.* 16: 1990-2001.
- Farkas, R.H., et al. 1997. Neurotensin and dopamine D2 activation oppositely regulate the same  $K^+$  conductance in rat midbrain dopaminergic neurons. *Neurosci. Letts.* 231: 21-24.
- Lei, Q., et al. 2000. Activation and inhibition of G protein-coupled inwardly rectifying potassium (KIR3) channels by G protein by subunits. *Proc. Natl. Acad. Sci.* 97: 9771-9776.
- Rogalski, S.L., et al. 2000. TrkB activation by brain-derived neurotrophic factor inhibits the G protein-gated inward rectifier KIR3 by tyrosine phosphorylation of the channel. *J. Biol. Chem.* 275: 25082-25088.

## CHROMOSOMAL LOCATION

Genetic locus: KCNJ13 (human) mapping to 2q37.1; Kcnj13 (mouse) mapping to 1 D.

## SOURCE

KIR7.1 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of KIR7.1 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-22438 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

KIR7.1 (N-16) is recommended for detection of KIR7.1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

KIR7.1 (N-16) is also recommended for detection of KIR7.1 in additional species, including equine and canine.

Suitable for use as control antibody for KIR7.1 siRNA (h): sc-42630, KIR7.1 siRNA (m): sc-155912, KIR7.1 shRNA Plasmid (h): sc-42630-SH, KIR7.1 shRNA Plasmid (m): sc-155912-SH, KIR7.1 shRNA (h) Lentiviral Particles: sc-42630-V and KIR7.1 shRNA (m) Lentiviral Particles: sc-155912-V.

Molecular Weight of KIR7.1: 67 kDa.

Positive Controls: mouse cerebellum extract: sc-2403 or rat cerebellum extract: sc-2398.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## SELECT PRODUCT CITATIONS

- Yang, D., et al. 2008. Expression of KIR7.1 and a novel KIR7.1 splice variant in native human retinal pigment epithelium. *Exp. Eye Res.* 86: 81-91.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfation  
Guaranteed

Try **KIR7.1 (C-12): sc-398810**, our highly recommended monoclonal alternative to KIR7.1 (N-16).