



Urease β (bN-19): sc-22446

BACKGROUND

Helicobacter pylori is present in the stomachs of at least half of the world's population. Although gastric adenocarcinoma is associated with the presence of *H. pylori* in the stomach, only a small fraction of colonized individuals develop this common malignancy. *H. pylori* urease, an enzyme that generates ammonia and is present within the lamina propria of colonized individuals, binds to class II major histocompatibility complex (MHC) molecules on the surfaces of gastric epithelial cells *in vitro*. In addition, Urease, which may be toxic to gastric epithelial cells, may also induce apoptosis. Specifically, Urease plays a crucial role in the development of ulcers in the duodenum by accelerating apoptosis in the antral mucosa. Ammonia accelerates TNF α cytokine-induced apoptosis, while ammonia or urease alone are unable to induce apoptosis. Urease exists as two forms, Urease α (UreA) and Urease β (UreB).

REFERENCES

- Smoot, D.T. 1997. How does *Helicobacter pylori* cause mucosal damage? Direct mechanisms. *Gastroenterology* 113: S31-34; discussion S50.
- Kohda, K., Tanaka, K., Aiba, Y., Yasuda, M., Miwa, T. and Koga, Y. 1999. Role of apoptosis induced by *Helicobacter pylori* infection in the development of duodenal ulcer. *Gut* 44: 456-462.
- Fan, X., Gunasena, H., Cheng, Z., Espejo, R., Crowe, S.E., Ernst, P.B. and Reyes, V.E. 2000. *Helicobacter pylori* urease binds to class II MHC on gastric epithelial cells and induces their apoptosis. *J. Immunol.* 165: 1918-1924.
- Igarashi, M., Kitada, Y., Yoshiyama, H., Takagi, A., Miwa, T. and Koga, Y. 2001. Ammonia as an accelerator of tumor necrosis factor α -induced apoptosis of gastric epithelial cells in *Helicobacter pylori* infection. *Infect. Immunol.* 69: 816-821.
- Kumagai, T., Yan, J., Graham, D.Y., Tozuka, M., Okimura, Y., Ikeno, T., Sugiyama, A., Katsuyama, T. and Ota, H. 2001. Serum immunoglobulin G immune response to *Helicobacter pylori* antigens in Mongolian gerbils. *J. Clin. Microbiol.* 39: 1283-1288.
- Peek, R.M., Jr. and Blaser, M.J. 2002. *Helicobacter pylori* and gastrointestinal tract adenocarcinomas. *Nat. Rev. Cancer* 2: 28-37.
- Lock, R.A., Coombs, G.W., McWilliams, T.M., Pearman, J.W., Grubb, W.B., Melrose, G.J. and Forbes, G.M. 2002. Proteome analysis of highly immunoreactive proteins of *Helicobacter pylori*. *Helicobacter* 7: 175-182.

SOURCE

Urease β (bN-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Urease β of *H. pylori* origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-22446 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Urease β (bN-19) is recommended for detection of Urease β of *H. pylori* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Urease β : 61 kDa.

Positive Controls: *E. coli* extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.