

NCK2 (V-14): sc-22508

BACKGROUND

The NCK family of SH2/SH3 adaptor proteins consists of two members, NCK1 (NCK α) and NCK2 (NCK β), which couple tyrosine kinase signaling, including the EGF and PDGF receptor-pathways, to downstream signaling proteins. Specifically, overexpression of NCK1 in NIH/3T3 cells decreases DNA synthesis stimulated by EGF. Furthermore, the SH2 domain of NCK2 inhibits EGF- and PDGF-induced DNA synthesis. The SH3 domain of NCK binds a proline-rich domain on PAK, a known actin cytoskeleton regulator. The NCK protein thus mediates the interaction between PAK and RAC. The NCK2 protein binds human PDGFR- β (Tyr 1009); overexpression of NCK2 inhibits PDGF-induced membrane ruffling and lamellipod formation. Various growth factor receptors, cell surface antigens and adhesion molecules phosphorylate mammalian NCK1 and NCK2. The human NCK1 and NCK2 genes map to chromosomes 3q21 and 2q12.2, respectively.

REFERENCES

1. Park, D., et al. 1992. Phosphorylation of NCK in response to a variety of receptors, phorbol myristate acetate, and cyclic AMP. *Mol. Cell. Biol.* 12: 5816-5823.
2. Huebner, K., et al. 1994. Chromosome locations of genes encoding human signal transduction adapter proteins, NCK, SHC1, and GRB2. *Genomics* 22: 281-287.
3. Chen, M., et al. 1998. Identification of NCK family genes, chromosomal localization, expression, and signaling specificity. *J. Biol. Chem.* 273: 25171-25178.
4. Chen, M., et al. 2000. NCK β adapter regulates actin polymerization in NIH/3T3 fibroblasts in response to platelet-derived growth factor $\beta\beta$. *Mol. Cell. Biol.* 20: 7867-7880.
5. Buday, L., et al. 2002. The NCK family of adapter proteins. Regulators of actin cytoskeleton. *Cell. Signal.* 14: 723-731.
6. Kiosses, W.B., et al. 2002. A dominant-negative p65 PAK peptide inhibits angiogenesis. *Circ. Res.* 90: 697-702.
7. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604930. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: NCK2 (human) mapping to 2q12.2; Nck2 (mouse) mapping to 1 B.

SOURCE

NCK2 (V-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of NCK2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-22508 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NCK2 (V-14) is recommended for detection of NCK2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NCK2 (V-14) is also recommended for detection of NCK2 in additional species, including equine and canine.

Suitable for use as control antibody for NCK2 siRNA (h): sc-36018, NCK2 siRNA (m): sc-40969, NCK2 shRNA Plasmid (h): sc-36018-SH, NCK2 shRNA Plasmid (m): sc-40969-SH, NCK2 shRNA (h) Lentiviral Particles: sc-36018-V and NCK2 shRNA (m) Lentiviral Particles: sc-40969-V.

Molecular Weight of NCK2: 47 kDa.

Positive Controls: U-937 cell lysate: sc-2239.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **NCK2 (8.8): sc-20020**, our highly recommended monoclonal alternative to NCK2 (V-14).