

# p- $\alpha$ B-crystallin (mSer 59)-R: sc-22510-R

## BACKGROUND

Crystallins are the major proteins of the vertebrate eye lens, where they maintain the transparency and refractive index of the lens. Crystallins are divided into  $\alpha$ ,  $\beta$ , and  $\gamma$  families, and the  $\beta$  and  $\gamma$ -crystallins also compose a superfamily. Crystallins usually contain seven distinctive protein regions, including four homologous motifs, a connecting peptide, and N- and C-terminal extensions.  $\alpha$ -crystallins consist of three gene products,  $\alpha$ A,  $\alpha$ B, and  $\alpha$ C-crystallin, which are members of the small heat shock protein family (HSP20). They are induced by heat shock, and act as molecular chaperones by holding denatured proteins in large soluble aggregates. However, unlike other molecular chaperones,  $\alpha$ -crystallins do not renature these proteins.  $\alpha$ A-crystallin is preferentially restricted to the lens. Defects in the gene encoding  $\alpha$ A-crystallin cause autosomal dominant congenital cataract (ADCC). The human  $\alpha$ B-crystallin gene encodes a protein that is present in many tissues, including lens, heart, and skeletal muscle. Elevated expression of  $\alpha$ B-crystallin is associated with many neurological diseases, and a missense mutation in this gene has cosegregated in a family with a desmin-related myopathy. The p38 MAPK substrate, MAPKAPK-2 phosphorylates  $\alpha$ B-crystallin on Ser 59.

## CHROMOSOMAL LOCATION

Genetic locus: CRYAB (human) mapping to 11q23.1; Cryab (mouse) mapping to 9 A5.3.

## SOURCE

p- $\alpha$ B-crystallin (mSer 59)-R is a rabbit polyclonal antibody raised against a short amino acid sequence containing Ser 59 phosphorylated  $\alpha$ B-crystallin of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-22510 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

p- $\alpha$ B-crystallin (mSer 59)-R is recommended for detection of Ser 59 phosphorylated  $\alpha$ B-crystallin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000). p- $\alpha$ B-crystallin (mSer 59)-R is also recommended for detection of correspondingly phosphorylated  $\alpha$ B-crystallin in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for  $\alpha$ B-crystallin siRNA (h): sc-40432,  $\alpha$ B-crystallin siRNA (m): sc-40433,  $\alpha$ B-crystallin shRNA Plasmid (h): sc-40432-SH,  $\alpha$ B-crystallin shRNA Plasmid (m): sc-40433-SH,  $\alpha$ B-crystallin shRNA (h) Lentiviral Particles: sc-40432-V and  $\alpha$ B-crystallin shRNA (m) Lentiviral Particles: sc-40433-V.

Molecular Weight of p- $\alpha$ B-crystallin: 20 kDa.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Western Blotting Luminol Reagent: sc-2048 and Lambda Phosphatase: sc-200312A. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## DATA



Western blot analysis of  $\alpha$ B-crystallin phosphorylation in untreated (A, C) and lambda protein phosphatase (sc-200312A) treated (B, D) rat kidney tissue extract. Antibodies tested include p- $\alpha$ B-crystallin (mSer 59)-R: sc-22510-R (A, B) and  $\alpha$ B-crystallin (F-10): sc-137129 (C, D).



Western blot analysis of  $\alpha$ B-crystallin phosphorylation in untreated (A, C) and lambda protein phosphatase (sc-200312A) treated (B, D) rat heart tissue extract. Antibodies tested include p- $\alpha$ B-crystallin (mSer 59)-R: sc-22510-R (A, B) and  $\alpha$ B-crystallin (F-10): sc-137129 (C, D).

## SELECT PRODUCT CITATIONS

- Schmeer, C., et al. 2008. Statins modulate heat shock protein expression and enhance retinal ganglion cell survival after transient retinal ischemia/reperfusion *in vivo*. Invest. Ophthalmol. Vis. Sci. 49: 4971-4981.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **p- $\alpha$ B-crystallin (F-1): sc-365884**, our highly recommended monoclonal alternative to p- $\alpha$ B-crystallin (mSer 59).