

E2F-3 (H-110): sc-22822

BACKGROUND

The human retinoblastoma gene product appears to play an important role in the negative regulation of cell proliferation. Functional inactivation of Rb can be mediated either through mutation or as a consequence of interaction with DNA tumor virus-encoded proteins. Of all the Rb associations described to date, the identification of a complex between Rb and the transcription factor E2F most directly implicates Rb in regulation of cell proliferation. E2F was originally identified through its role in transcriptional activation of the adenovirus E2 promoter. Sequences homologous to the E2F binding site have been found upstream of a number of genes that encode proteins with putative functions in the G₁ and S phases of the cell cycle. E2F-1 is a member of a broader family of transcription regulators including E2F-2, E2F-3, E2F-4, E2F-5 and E2F-6, each of which forms heterodimers with a second protein, DP-1, forming an "active" E2F transcriptional regulatory complex.

REFERENCES

1. Chellappan, S., et al. 1991. The E2F transcription factor is a cellular target for the Rb protein. *Cell* 65: 1053-1061.
2. Chittenden, T., et al. 1991. The T/E1A-binding domain of the retinoblastoma product can interact selectively with a sequence-specific DNA-binding protein. *Cell* 65: 1073-1082.
3. Helin, K., et al. 1992. A cDNA encoding a pRb-binding protein with properties of the transcription factor E2F. *Cell* 70: 337-350.
4. Helin, K., et al. 1993. Heterodimerization of the transcription factors E2F-1 and DP-1 leads to cooperative transactivation. *Genes Dev.* 7: 1850-1861.
5. Krek, W., et al. 1993. Binding to DNA and the retinoblastoma gene product promoted by complex formation of different E2F family members. *Science* 262: 1557-1560.
6. Ginsberg, D., et al. 1994. E2F-4, a new member of the E2F transcription factor family, interacts with p107. *Genes Dev.* 8: 2665-2679.

CHROMOSOMAL LOCATION

Genetic locus: E2F3 (human) mapping to 6p22.3; E2f3 (mouse) mapping to 13 A3.2.

SOURCE

E2F-3 (H-110) is a rabbit polyclonal antibody raised against amino acids 1-110 of E2F-3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-22822 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

E2F-3 (H-110) is recommended for detection of E2F-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

E2F-3 (H-110) is also recommended for detection of E2F-3 in additional species, including bovine and equine.


Suitable for use as control antibody for E2F-3 siRNA (h): sc-37817, E2F-3 siRNA (m): sc-37818, E2F-3 shRNA Plasmid (h): sc-37817-SH, E2F-3 shRNA Plasmid (m): sc-37818-SH, E2F-3 shRNA (h) Lentiviral Particles: sc-37817-V and E2F-3 shRNA (m) Lentiviral Particles: sc-37818-V.

E2F-3 (H-110) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of E2F-3: 55 kDa.

Positive Controls: Y79 nuclear extract: sc-2126, E2F-3 (h): 293 Lysate: sc-113198 or K-562 nuclear extract: sc-2130.

DATA



E2F-3 (H-110): sc-22822. Western blot analysis E2F-3 expression in K-562 (A) and Y79 (B) whole cell lysates and K-562 (C) and Y79 (D) nuclear extracts.



E2F-3 (H-110): sc-22822. Western blot analysis of E2F-3 expression in non-transfected 293: sc-110760 (A), human E2F-3 transfected 293: sc-113198 (B) and HeLa (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **E2F-3 (PG30): sc-56665** or **E2F-3 (D-2): sc-28308**, our highly recommended monoclonal alternatives to E2F-3 (H-110).