

Myf-5 (H-50): sc-22825

BACKGROUND

Differentiation of myogenic cells is regulated by multiple positively and negatively acting factors. One well characterized family of helix-loop-helix (HLH) proteins known to play an important role in the regulation of muscle cell development includes MyoD, myogenin, Myf-5 and Myf-6 (also designated MRF-4 or herculin). Of interest, most muscle cells express either MyoD or Myf-5 in the committed state, but when induced to differentiate, all turn on expression of myogenin. MyoD transcription factors form heterodimers with products of a more widely expressed family of bHLH genes, the E family, which consists of at least three distinct genes: E2A, IF2 and HEB. MyoD-E heterodimers bind avidly to consensus (CANNTG) E box target sites that are functionally important elements in the upstream regulatory sequences of many muscle-specific terminal differentiation genes.

REFERENCES

1. Braun, T., et al. 1989. A novel human muscle factor related to but distinct from Myo D1 induces myogenic conversion in 10T1/2 fibroblasts. *EMBO J.* 8: 701-709.
2. Rhodes, S.J., et al. 1989. Identification of MRF4: a new member of the muscle regulatory factor gene family. *Genes Dev.* 3: 2050-2061.
3. Wright, W.E., et al. 1989. Myogenin, a factor regulating myogenesis, has a domain homologous to Myo D. *Cell* 56: 607-617.
4. Miner, J.H., et al. 1990. Herculin, a fourth member of the Myo D family of myogenic regulatory genes. *Proc. Natl. Acad. Sci. USA* 87: 1089-1093.
5. Braun, T., et al. 1990. Myf-6, a new member of the human gene family of myogenic determination factors: evidence for a gene cluster on chromosome 12. *EMBO J.* 9: 821-831.

CHROMOSOMAL LOCATION

Genetic locus: MYF5 (human) mapping to 12q21.31; Myf5 (mouse) mapping to 10 D1.

SOURCE

Myf-5 (H-50) is a rabbit polyclonal antibody raised against amino acids 206-255 mapping at the C-terminus of Myf-5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-22825 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Myf-5 (H-50) is recommended for detection of Myf-5 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Myf-5 siRNA (h): sc-35988, Myf-5 siRNA (m): sc-35989, Myf-5 shRNA Plasmid (h): sc-35988-SH, Myf-5 shRNA Plasmid (m): sc-35989-SH, Myf-5 shRNA (h) Lentiviral Particles: sc-35988-V and Myf-5 shRNA (m) Lentiviral Particles: sc-35989-V.


Myf-5 (H-50) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of Myf-5: 32 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



Myf-5 (H-50): sc-22825. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic staining of myocytes.

SELECT PRODUCT CITATIONS

1. Hanssen, K.E., et al. 2012. The effect of strength training volume on satellite cells, myogenic regulatory factors, and growth factors. *Scand. J. Med. Sci. Sports.* E-published.