

VEGI (P-18): sc-23186

BACKGROUND

Vascular endothelial cell growth inhibitor (VEGI), also known as TNFRSF15 or TL1, a member of the TNF superfamily, has a signaling pathway similar to TNF and is most likely a multifunctional cytokine. VEGI is found in brain, reproductive and late developmental stage embryonic tissues and expressed predominantly in endothelial cells. It is an angiogenesis inhibitor of the TNF family and functions in part by directly inhibiting endothelial cell proliferation. VEGI may act as an autocrine factor to induce apoptosis in endothelial cells via activation of multiple signaling pathways, including stress protein kinases as well as certain caspases.

REFERENCES

1. Tan, K.B., et al. 1997. Characterization of a novel TNF-like ligand and recently described TNF ligand and TNF receptor superfamily genes and their constitutive and inducible expression in hematopoietic and non-hematopoietic cells. *Gene* 204: 35-46.
2. Haridas, V., et al. 1999. VEGI, a new member of the TNF family activates nuclear factor κ B and c-Jun N-terminal kinase and modulates cell growth. *Oncogene* 18: 6496-6504.
3. Hu, S., et al. 1999. Characterization of TNFRSF19, a novel member of the tumor necrosis factor receptor superfamily. *Genomics* 62: 103-107.
4. Zhai, Y., et al. 1999. Inhibition of angiogenesis and breast cancer xenograft tumor growth by VEGI, a novel cytokine of the TNF superfamily. *Int. J. Cancer* 82: 131-136.
5. Zhai, Y., et al. 1999. VEGI, a novel cytokine of the tumor necrosis factor family, is an angiogenesis inhibitor that suppresses the growth of colon carcinomas *in vivo*. *FASEB J.* 13: 181-189.
6. Yue, T.L., et al. 1999. TL1, a novel tumor necrosis factor-like cytokine, induces apoptosis in endothelial cells. Involvement of activation of stress protein kinases (stress-activated protein kinase and p38 mitogen-activated protein kinase) and caspase-3-like protease. *J. Biol. Chem.* 274: 1479-1486.

CHROMOSOMAL LOCATION

Genetic locus: TNFSF15 (human) mapping to 9q32; Tnfsf15 (mouse) mapping to 4 C1.

SOURCE

VEGI (P-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of VEGI of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-23186 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

VEGI (P-18) is recommended for detection of VEGI of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

VEGI (P-18) is also recommended for detection of VEGI in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for VEGI siRNA (h): sc-39846, VEGI siRNA (m): sc-39847, VEGI shRNA Plasmid (h): sc-39846-SH, VEGI shRNA Plasmid (m): sc-39847-SH, VEGI shRNA (h) Lentiviral Particles: sc-39846-V and VEGI shRNA (m) Lentiviral Particles: sc-39847-V.

Molecular Weight of VEGI: 22 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



VEGI (P-18): sc-23186. Western blot analysis of human recombinant VEGI.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **VEGI (1A3): sc-53975**, our highly recommended monoclonal alternative to VEGI (P-18).