

p44S10 (K-16): sc-23208



The Power to Question

BACKGROUND

In eukaryotic cells, selective breakdown of cellular proteins is ensured by two distinct pathways, ubiquitination and degradation by the 26S proteasome. At specific stages of development, embryo- and tissue-specific components of the 26S proteasome are formed by developmentally regulated alternative splicing, including Rpn10a through Rpn10e (also designated pUb-R2 through pUb-R5). The pUb-R2 subunit, originally identified as S5a, is ubiquitously expressed and may perform proteolysis constitutively in a wide variety of cells. p44S10 is a highly conserved proteasome regulatory subunit that is expressed in heart, liver, skeletal muscle and pancreas. In addition to normal tissue expression, p44S10 is also expressed in several melanoma cell lines, such as MCF-7, 451Lu and WM164. Since forced expression of p44S10 in radial growth phase melanoma cells results in an increase in cellular proliferation, p44S10 may represent a potential link between regulation of proteasome activity and tumor cell proliferation *in vivo*.

REFERENCES

- Lonroth, I. and Lange, S. 1986. Purification and characterization of the antisecretory factor: a protein in the central nervous system and in the gut which inhibits intestinal hypersecretion induced by cholera toxin. *Biochim. Biophys. Acta* 883: 138-144.
- Johansson, E., Lonroth, I., Lange, S., Jonson, I., Jennische, E. and Lonroth, C. 1995. Molecular cloning and expression of a pituitary gland protein modulating intestinal fluid secretion. *J. Biol. Chem.* 270: 20615-20620.
- Coux, O., Tanaka, K. and Goldberg, A.L. 1996. Structure and functions of the 20S and 26S proteasomes. *Annu. Rev. Biochem.* 65: 801-847.
- Voges, D., Zwickl, P. and Baumeister, W. 1999. The 26S proteasome: a molecular machine designed for controlled proteolysis. *Annu. Rev. Biochem.* 68: 1015-1068.
- Kawahara, H., Kasahara, M., Nishiyama, A., Ohsumi, K., Goto, T., Kishimoto, T., Saeki, Y., Yokosawa, H., Shimbara, N., Murata, S., Chiba, T., Suzuki, K. and Tanaka, K. 2000. Developmentally regulated, alternative splicing of the Rpn10 gene generates multiple forms of 26S proteasomes. *EMBO J.* 19: 4144-4153.
- Ren, S., Smith, M.J., Louro, I.D., McKie-Bell, P., Bani, M.R., Wagner, M., Zochodne, B., Redden, D.T., Grizzle, W.E., Wang, N.D., Smith, D.I., Herbst, R.A., Bardenheuer, W., Opalka, B., Schutte, J., Trent, J.M., Ben-David, Y. and Ruppert, J.M. 2000. The p44S10 locus, encoding a subunit of the proteasome regulatory particle, is amplified during progression of cutaneous malignant melanoma. *Oncogene* 19: 1419-1427.

CHROMOSOMAL LOCATION

Genetic locus: PSMD6 (human) mapping to 3p14.1; Psmd6 (mouse) mapping to 14 A1.

SOURCE

p44S10 (K-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of p44S10 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-23208 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

p44S10 (K-16) is recommended for detection of p44S10 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

p44S10 (K-16) is also recommended for detection of p44S10 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for p44S10 siRNA (h): sc-41383, p44S10 siRNA (m): sc-41384, p44S10 shRNA Plasmid (h): sc-41383-SH, p44S10 shRNA Plasmid (m): sc-41384-SH, p44S10 shRNA (h) Lentiviral Particles: sc-41383-V and p44S10 shRNA (m) Lentiviral Particles: sc-41384-V.

Molecular Weight of p44S10: 44 kDa.

Positive Controls: Rat heart extract: sc-2393, A-375 cell lysate: sc-3811 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **p44S10 (E-12): sc-393580** or **p44S10 (H-5): sc-398745**, our highly recommended monoclonal alternatives to p44S10 (K-16).