

# TRPV5 (C-15): sc-23376

## BACKGROUND

Transient receptor potential (TRP) proteins are cation-sensitive channels that modulate a myriad of cellular functions, including temperature sensation and vasoregulation. Transcribed from a gene adjacent to VR-1, the thermal-sensitive, capsaicin-insensitive TRPV3 is expressed at warm temperatures; expression increases in response to noxious temperatures. Human TRPV3 is expressed in skin, tongue, dorsal root ganglion, trigeminal ganglion, spinal cord and brain. In addition, TRPV3 is co-expressed in dorsal root ganglion neurons with VR-1. TRPV3 associates with VR-1 and may modulate VR-1 activity. The 729 amino acid TRPV5 (ECAC1) protein comprises six trans-membrane domains, multiple potential phosphorylation sites, an N-linked glycosylation site and three ankyrin repeat regions. It is abundantly expressed in kidney, jejunum and pancreas and at lower levels in testis, prostate, placenta, brain, colon and rectum. TRPV5 controls the rate-limiting step of vitamin D3-regulated Ca<sup>2+</sup> reabsorption in kidney and intestine; the 5'-flanking region of TRPV5 contains four putative vitamin D3-responsive elements.

## REFERENCES

- Muller, D., Hoenderop, J.G., Merckx, G.F., van Os, C.H. and Bindels, R.J. 2000. Gene structure and chromosomal mapping of human epithelial calcium channel. *Biochem. Biophys. Res. Commun.* 275: 47-52.
- Smith, G.D., Gunthorpe, M.J., Kelsell, R.E., Hayes, P.D., Reilly, P., Facer, P., Wright, J.E., Jerman, J.C., Walhin, J.p., Ooi, L., Egerton, J., Charles, K.J., Smart, D., Randall, A.D., Anand, P. and Davis, J.B. 2000. TRPV3 is a temperature-sensitive vanilloid receptor-like protein. *Nature* 418: 186-190.
- Peier, A.M., Reeve, A.J., Andersson, D.A., Moqrich, A., Earley, T.J., Hegarden, A.C., Story, G.M., Colley, S., Hogenesch, J.B., McIntyre, P., Bevan, S. and Patapoutian, A. 2002. A heat-sensitive TRP channel expressed in keratinocytes. *Science* 296: 2046-2049.
- Xu, H., Ramsey, L.S., Kotecha, S.A., Moran, M.M., Chong, J.A., Lawson, D., Ge, P., Lilly, J., Silo-Santiago, I., Xie, Y., DiStefano, P.S., Curtis, R. and Clapham, D.E. 2002. TRPV3 is a calcium-permeable temperature-sensitive cation channel. *Nature* 418: 181-186.

## CHROMOSOMAL LOCATION

Genetic locus: TRPV5 (human) mapping to 7q31.1; Trpv5 (mouse) mapping to 6 B2.1.

## SOURCE

TRPV5 (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of TRPV5 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-23376 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

TRPV5 (C-15) is recommended for detection of TRPV5 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TRPV5 siRNA (h): sc-42676, TRPV5 shRNA Plasmid (h): sc-42676-SH and TRPV5 shRNA (h) Lentiviral Particles: sc-42676-V.

Molecular Weight of TRPV5 core: 75 kDa.

Molecular Weight of glycosylated TRPV5: 85-100 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **TRPV5 (B-6): sc-398853**, our highly recommended monoclonal alternative to TRPV5 (C-15).