

Aiolos (S-21): sc-23586

BACKGROUND

Ikaros family members, including Ikaros and Helios, are nuclear factors that colocalize with DNA replication machinery components in higher-order chromatin structures and respond to signaling events, such as T cell activation. Helios and Ikaros bind to similar DNA sequences and they function as hematopoietic-specific transcription factors. Members of the Ikaros family contain zinc-finger domains that are involved in DNA-binding and in the formation of homodimers and heterodimers between Ikaros family members. Aiolos, also known as zinc finger protein Ikaros 3 or ZNF1A3, is a 509 amino acid nuclear protein. Aiolos plays an important role in lymphocyte differentiation regulation and, via this role, mutated Aiolos is implicated in leukemogenesis. Expressed in most tissues, Aiolos is predominantly found in spleen, thymus and peripheral blood leukocytes. Aiolos contains six C₂H₂-type zinc fingers, a motif commonly involved in nucleotide binding. Aiolos interacts with Ikaros family members, including Eos and Pegasus.

REFERENCES

1. Wang, J.H., et al. 1998. Aiolos regulates B cell activation and maturation to effector state. *Immunity* 9: 543-553.
2. Hosokawa, Y., et al. 1999. Human aiolos, an ikaros-related zinc finger DNA binding protein: cDNA cloning, tissue expression pattern, and chromosomal mapping. *Genomics* 61: 326-329.
3. Perdomo, J., et al. 2000. Eos and pegasus, two members of the Ikaros family of proteins with distinct DNA binding activities. *J. Biol. Chem.* 275: 38347-38354.
4. Liippo, J., et al. 2001. Both normal and leukemic B lymphocytes express multiple isoforms of the human Aiolos gene. *Eur. J. Immunol.* 31: 3469-3474.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606221. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Sun, J., et al. 2003. Lack of the transcriptional coactivator OBF-1 prevents the development of systemic lupus erythematosus-like phenotypes in Aiolos mutant mice. *J. Immunol.* 170: 1699-1706.

CHROMOSOMAL LOCATION

Genetic locus: Ikzf3 (mouse) mapping to 11 D.

SOURCE

Aiolos (S-21) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Aiolos of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-23586 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-23586 X, 200 µg/0.1 ml.

APPLICATIONS

Aiolos (S-21) is recommended for detection of Aiolos of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Aiolos siRNA (m): sc-38006, Aiolos shRNA Plasmid (m): sc-38006-SH and Aiolos shRNA (m) Lentiviral Particles: sc-38006-V.

Aiolos (S-21) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of Aiolos: 58 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Aiolos (3H5-G7): sc-293421**, our highly recommended monoclonal alternative to Aiolos (S-21).