

# MAAI (N-20): sc-23693

## BACKGROUND

In humans, maleylacetoacetate isomerase (MAAI, also known as GSTZ1-1) catalyzes the conversion of maleylacetoacetate to fumarylacetoacetate, the fifth step in the phenylalanine/phenylacetate degradation pathway. Deficiencies in other steps of this pathway cause metabolic diseases, including type I tyrosinemia and phenylketonuria. The glutathione S-transferases (GSTs) are found in all aerobic organisms and catalyze the conjugation of glutathione to a wide variety of electrophilic substrates. By sequence alignment and phylogenetic analysis, a new subgroup of GST-like proteins from human, *C. elegans*, and carnation were identified. Human MAAI is 38% and 49% identical to the carnation and *C. elegans* proteins, respectively. Recombinant human MAAI is a dimer. The enzyme exhibits limited activity with known GST substrates. Western blot analysis indicates that MAAI is most abundant in liver, with lower levels detected in skeletal muscle and brain. The gene which encodes MAAI maps to human chromosome 14q24.3.

## REFERENCES

- Berger, R., et al. 1988. Tyrosinemia type Ib caused by maleylacetoacetate isomerase deficiency: a new enzyme defect. *Pediatr. Res.* 23: 328A.
- Board, P.G., et al. 1997.  $\zeta$ , a novel class of glutathione transferases in a range of species from plants to humans. *Biochem. J.* 328: 929-935.
- Blackburn, A.C., et al. 1998. Characterization and chromosome location of the gene GSTZ1 encoding the human  $\zeta$  class glutathione transferase and maleylacetoacetate isomerase. *Cytogenet. Cell Genet.* 83: 109-114.
- Fernandez-Canon, J.M., et al. 1998. Characterization of a fungal maleylacetoacetate isomerase gene and identification of its human homologue. *J. Biol. Chem.* 273: 329-337.
- Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603758. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: GSTZ1 (human) mapping to 14q24.3; Gstz1 (mouse) mapping to 12 D2.

## SOURCE

MAAI (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of MAAI of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-23693 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

MAAI (N-20) is recommended for detection of MAAI of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MAAI (N-20) is also recommended for detection of MAAI in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for MAAI siRNA (h): sc-40729, MAAI siRNA (m): sc-40730, MAAI shRNA Plasmid (h): sc-40729-SH, MAAI shRNA Plasmid (m): sc-40730-SH, MAAI shRNA (h) Lentiviral Particles: sc-40729-V and MAAI shRNA (m) Lentiviral Particles: sc-40730-V.


Molecular Weight of MAAI: 24.1 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227 or mouse liver extract: sc-2256.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz<sup>™</sup>: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



MAAI (N-20): sc-23693. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing cytoplasmic staining of hepatocytes.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.