NPY5-R (C-20): sc-23845



The Power to Overtin

BACKGROUND

Pancreatic polypeptide (PP), neuropeptide Y (NPY), and peptide YY (PYY) are related 36-amino acid hormones. A number of structurally related receptors for these peptides have been isolated, NPY1-R, NPY2-R, NPY3-R, NPY4-R, NPY5-R, and NPY6-R. NPY4-R is expressed in several human tissues, including brain, coronary artery, and ileum. NPY4-R maps to human chromosome 10q11.22. NPY-5R, isolated from rat hypothalamus, encodes a 456-amino acid protein with less than 35% overall identity to known Y-type receptors. The human NPY-5R sequence is nearly identical to, but in the opposite orientation from, that of the human NPY-1R sequence. NPY5-R localizes to the paraventricular hypothalamic nucleus, the lateral hypothalamus, and other locations consistent with a role in the control of feeding behavior. The gene which encodes NPY-5R maps to human chromosome 4q32.2. NPY-6R is abundantly expressed in human heart and skeletal muscle and the gene which encodes NPY-6R maps to human chromosome 5q31.

REFERENCES

- Bard, J.A., et al. 1995. Cloning and functional expression of a human Y4 subtype receptor for pancreatic polypeptide, neuropeptide Y, and peptide YY. J. Biol. Chem. 270: 26762-26765.
- Gerald, C., et al. 1996. A receptor subtype involved in neuropeptide-Yinduced food intake. Nature 382: 168-171.
- Hu, Y., et al. 1996. Identification of a novel hypothalamic neuropeptide Y receptor associated with feeding behavior. J. Biol. Chem. 271: 26315-26319.
- 4. Matsumoto, M., et al. 1996. Inactivation of a novel neuropeptide Y/peptide YY receptor gene in primate species. J. Biol. Chem. 271: 27217-27220.
- Herzog, H., et al. 1997. Overlapping gene structure of the human neuropeptide Y receptor subtypes Y1 and Y5 suggests coordinate transcriptional regulation. Genomics 41: 315-319.
- 6. Lutz, C.M., et al. 1997. Neuropeptide Y receptor genes mapped in human and mouse: receptors with high affinity for pancreatic polypeptide are not clustered with receptors specific for neuropeptide Y and peptide YY. Genomics 46: 287-290.
- Darby, K., et al. 1997. Assignment of the Y-4 receptor gene (PPYR1) to human chromosome 10q11.2 and mouse chromosome 14. Genomics 46: 513-515.

CHROMOSOMAL LOCATION

Genetic locus: NPY5R (human) mapping to 4q32.2; Npy5r (mouse) mapping to 8 B3.2.

SOURCE

NPY5-R (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of NPY5-R of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-23845 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NPY5-R (C-20) is recommended for detection of NPY5-R of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


NPY5-R (C-20) is also recommended for detection of NPY5-R in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for NPY5-R siRNA (h): sc-42105, NPY5-R siRNA (m): sc-42106, NPY5-R shRNA Plasmid (h): sc-42105-SH, NPY5-R shRNA Plasmid (m): sc-42106-SH, NPY5-R shRNA (h) Lentiviral Particles: sc-42105-V and NPY5-R shRNA (m) Lentiviral Particles: sc-42106-V.

Molecular Weight of NPY5-R: 57 kDa.

Positive Controls: MIA PaCa-2 cell lysate: sc-2285, Ramos cell lysate: sc-2216 or SK-N-SH cell lysate: sc-2410.

DATA



NPY5-R (C-20): sc-23845. Western blot analysis of NPY5-R expression in MIA PaCa-2 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS Satisfation Guaranteed

Try **NPY5-R (H-4): sc-137167**, our highly recommended monoclonal alternative to NPY5-R (C-20).