

# APC1 (P-20): sc-23963

## BACKGROUND

The anaphase-promoting complex (APC) is composed of more than ten subunits, including APC1, APC2, APC4, APC5, APC7, APC8, APC10, and APC11. The APC acts in a cell-cycle dependent manner to promote the separation of sister chromatids during the transition between metaphase and anaphase in mitosis. APC, or cyclosome, accomplishes this progression through the ubiquitination of mitotic cyclins and other regulatory proteins that are targeted for destruction during cell division. APC is phosphorylated, and thus activated, by protein kinases Cdk1/cyclin B and polo-like kinase (Plk). APC is under tight control by a number of regulatory factors, including CDC20, CDH1 and MAD2. Specifically, CDC20 and CDH1 directly bind to and activate APC's cyclin-ubiquitination activity. In contrast, MAD2 inhibits APC by forming a ternary complex with CDC20 and APC; thus preventing APC activation. APC1, previously referred to as Tsg24 in mice, is the largest of the APC subunits and is encoded by a gene mapping to 2q13.

## REFERENCES

1. Jorgensen, P.M., et al. 1998. A subunit of the anaphase-promoting complex is a centromere-associated protein in mammalian cells. *Mol. Cell. Biol.* 18: 468-476.
2. Page, A.M., et al. 1999. The anaphase-promoting complex: new subunits and regulators. *Annu. Rev. Biochem.* 68: 583-609.
3. Peters, J.M. 1999. Subunits and substrates of the anaphase-promoting complex. *Exp. Cell Res.* 248: 339-349.
4. Fang, G., et al. 1999. Control of mitotic transitions by the anaphase-promoting complex. *Phil. Trans. Royal Soc. London Biol. Sci.* 354: 1583-1590.
5. Jorgensen, P.M., et al. 2001. Characterisation of the human APC1, the largest subunit of the anaphase-promoting complex. *Gene* 262: 51-59.
6. Bolte, M., et al. 2002. Inhibition of APC-mediated proteolysis by the meiosis-specific protein kinase Ime2. *Proc. Natl. Acad. Sci. USA* 99: 4385-4390.
7. Golan, A., et al. 2002. The cyclin-ubiquitin ligase activity of cyclosome/APC is jointly activated by protein kinases Cdk1/cyclin B and Plk. *J. Biol. Chem.* 277: 15552-15557.

## CHROMOSOMAL LOCATION

Genetic locus: ANAPC1 (human) mapping to 2q13; Anapc1 (mouse) mapping to 2 F1.

## SOURCE

APC1 (P-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of APC1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-23963 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

APC1 (P-20) is recommended for detection of APC1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

APC1 (P-20) is also recommended for detection of APC1 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for APC1 siRNA (h): sc-37524, APC1 siRNA (m): sc-37525, APC1 shRNA Plasmid (h): sc-37524-SH, APC1 shRNA Plasmid (m): sc-37525-SH, APC1 shRNA (h) Lentiviral Particles: sc-37524-V and APC1 shRNA (m) Lentiviral Particles: sc-37525-V.

Molecular Weight of APC1: 243 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.