

# PRX (C-18): sc-23970

## BACKGROUND

The peroxiredoxin (PRX) family comprises six antioxidant proteins, PRX I, II, III, IV, V, and VI, which protect cells from reactive oxygen species (ROS) by preventing the metal-catalyzed oxidation of enzymes. The PRX proteins primarily utilize thioredoxin as the electron donor for antioxidation, although they are fairly promiscuous with regard to the hydroperoxide substrate. In addition to protection from ROS, peroxiredoxins are also involved in cell proliferation, differentiation, and gene expression. PRX I, III, IV, and VI show diffuse cytoplasmic localization, while PRX III and V exhibit distinct mitochondrial localization. The human PRX I gene encodes a protein that is expressed in several tissues, including liver, kidney, testis, lung, and the nervous system. The human PRX II and III genes map to chromosomes 13q12 and 10q25-q26, respectively. PRX II is expressed in testis, while PRX III shows expression in lung. PRX I, II and III are overexpressed in breast cancer and may be involved in its development or progression. Also, up-regulated protein levels of PRX I and II in Alzheimer's disease (AD) and Down Syndrome (DS) indicate the involvement of PRX I and II in their pathogenesis. The human PRX IV gene is abundantly expressed in many tissues. PRX IV exists as a precursor protein, which is only detected in testis, and a processed secreted form. PRX V also exists as two forms, designated long and short. Like PRX IV, the long form of PRX V is highly expressed in testis. The short form of PRX V is more widely expressed, with high expression in liver, kidney, heart and lung.

## REFERENCES

- Iwahara, S., et al. 1995. Purification, characterization, and cloning of a heme-binding protein (23 kDa) in rat liver cytosol. *Biochemistry* 34: 13398-13406.
- Butterfield, L.H., et al. 1999. From cytoprotection to tumor suppression: the multifactorial role of peroxiredoxins. *Antioxid. Redox Signal.* 1: 385-402.
- Noh, D.Y., et al. 2001. Over-expression of peroxiredoxin in human breast cancer. *Anticancer Res.* 21: 2085-2090.
- Kim, S.H., et al. 2001. Protein levels of human peroxiredoxin subtypes in brains of patients with Alzheimer's disease and Down syndrome. *J. Neural Transm. Suppl.* 223-235.
- Kinnula, V.L., et al. 2002. Cell specific expression of peroxiredoxins in human lung and pulmonary sarcoidosis. *Thorax* 57: 157-164.
- Hofmann, B., et al. 2002. Peroxiredoxins. *Biol. Chem.* 383: 347-364.

## SOURCE

PRX (C-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of PRX II of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-23970 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

PRX (C-18) is recommended for detection of PRX I, II, IV and, to a lesser extent, PRX III of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


PRX (C-18) is also recommended for detection of PRX I, II, IV and, to a lesser extent, PRX III in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for PRX siRNA (h): sc-37151, PRX siRNA (m): sc-37152, PRX shRNA Plasmid (h): sc-37151-SH, PRX shRNA Plasmid (m): sc-37152-SH, PRX shRNA (h) Lentiviral Particles: sc-37151-V and PRX shRNA (m) Lentiviral Particles: sc-37152-V.


Molecular Weight of PRX: 25 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206 or PRX II (h): 293T Lysate: sc-117298 or PRX II (m): 293T Lysate: sc-122808.

## DATA



PRX (C-18): sc-23970. Western blot analysis of PRX II expression in non-transfected: sc-117752 (A) and human PRX II transfected: sc-117298 (B) 293T whole cell lysates.



PRX (C-18): sc-23970. Western blot analysis of PRX II expression in non-transfected: sc-117752 (A) and mouse PRX II transfected: sc-122808 (B) 293T whole cell lysates.

## SELECT PRODUCT CITATIONS

- Yu, X., et al. 2006. The regulation of exosome secretion: a novel function of the p53 protein. *Cancer Res.* 66: 4795-4801.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **PRX (B-11): sc-137222** or **PRX (E-7): sc-271020**, our highly recommended monoclonal alternatives to PRX (C-18).