

COL5A3 (C-13): sc-240246

BACKGROUND

The extensive family of COL gene products (collagens) is composed of several chain types, including fibril-forming interstitial collagens (types I, II, III and V) and basement membrane collagens (type IV), each type containing multiple isoforms. Collagens are fibrous, extracellular matrix proteins with high tensile strength and are the major components of connective tissue, such as tendons and cartilage. All collagens contain a triple helix domain and frequently show lateral self-association in order to form complex connective tissues. Several collagens also play a role in cell adhesion, important for maintaining normal tissue architecture and function. COL5A3 (collagen α -3(V) chain) is a 1,745 amino acid extracellular matrix protein that belongs to the fibrillar collagen family and acts as a minor connective tissue. COL5A3 contains one TSP N-terminal (TSPN) domain, a fibrillar collagen NC1 domain and is encoded by a gene that maps to human chromosome 19p13.2.

REFERENCES

- Mann, K. 1992. Isolation of the α 3-chain of human type V collagen and characterization by partial sequencing. *Biol. Chem. Hoppe-Seyler* 373: 69-75.
- McCarthy, J.B., Vachhani, B. and Iida, J. 1996. Cell adhesion to collagenous matrices. *Biopolymers* 40: 371-381.
- Cremer, M.A., Rosloniec, E.F. and Kang, A.H. 1998. The cartilage collagens: a review of their structure, organization, and role in the pathogenesis of experimental arthritis in animals and in human rheumatic disease. *J. Mol. Med.* 76: 275-288.
- Alberio, L. and Dale, G.L. 1999. Review article: platelet-collagen interactions: membrane receptors and intracellular signalling pathways. *Eur. J. Clin. Invest.* 29: 1066-1076.
- Boskey, A.L., Wright, T.M. and Blank, R.D. 1999. Collagen and bone strength. *J. Bone Miner. Res.* 14: 330-335.
- Imamura, Y., Scott, I.C. and Greenspan, D.S. 2000. The pro- α 3(V) collagen chain. Complete primary structure, expression domains in adult and developing tissues, and comparison to the structures and expression domains of the other types V and XI procollagen chains. *J. Biol. Chem.* 275: 8749-8759.
- Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 120216. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: COL5A3 (human) mapping to 19p13.2; Col5a3 (mouse) mapping to 9 A3.

SOURCE

COL5A3 (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of Collagen α 3 Type V of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-240246 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

COL5A3 (C-13) is recommended for detection of Collagen α 3 Type V of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other COL5A family members.

COL5A3 (C-13) is also recommended for detection of Collagen α 3 Type V in additional species, including canine and porcine.

Suitable for use as control antibody for COL5A3 siRNA (h): sc-97621, COL5A3 siRNA (m): sc-142472, COL5A3 shRNA Plasmid (h): sc-97621-SH, COL5A3 shRNA Plasmid (m): sc-142472-SH, COL5A3 shRNA (h) Lentiviral Particles: sc-97621-V and COL5A3 shRNA (m) Lentiviral Particles: sc-142472-V.

Molecular Weight of COL5A3: 172 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.