

SERF1 (K-13): sc-240856

BACKGROUND

SERF1 (small EDRK-rich factor 1), also known as protein 4F5, SMAM1 (SMA modifier 1), SERF1A or SERF1B, is a 110 amino acid protein that exists as two alternatively spliced isoforms, designated isoform short and isoform long. Both SERF1 isoforms are expressed in spinal cord and central nervous system, but only isoform long is found in heart and skeletal muscle. The gene encoding SERF1 maps to human chromosome 5, which contains 181 million base pairs and comprises nearly 6% of the human genome. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

- Scharf, J.M., et al. 1996. The mouse region syntenic for human spinal muscular atrophy lies within the Lgn1 critical interval and contains multiple copies of Naip exon 5. *Genomics* 38: 405-417.
- Edwards, S.J., et al. 1997. The mutational spectrum in Treacher Collins syndrome reveals a predominance of mutations that create a premature-termination codon. *Am. J. Hum. Genet.* 60: 515-524.
- McDaniel, L.D., et al. 1997. Confirmation of homozygosity for a single nucleotide substitution mutation in a Cockayne syndrome patient using monoallelic mutation analysis in somatic cell hybrids. *Hum. Mutat.* 10: 317-321.
- Scharf, J.M., et al. 1998. Identification of a candidate modifying gene for spinal muscular atrophy by comparative genomics. *Nat. Genet.* 20: 83-86.
- Online Mendelian Inheritance in Man, OMIM[™]. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 603011. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Vera-Carbonell, A., et al. 2009. Characterization of a *de novo* complex chromosomal rearrangement in a patient with cri-du-chat and trisomy 5p syndromes. *Am. J. Med. Genet. A* 149A: 2513-2521.
- Ravandi, F., et al. 2009. Superior outcome with hypomethylating therapy in patients with acute myeloid leukemia and high-risk myelodysplastic syndrome and chromosome 5 and 7 abnormalities. *Cancer* 115: 5746-5751.
- Sazawal, S., et al. 2009. Haematological and molecular profile of acute myelogenous leukaemia in India. *Indian J. Med. Res.* 129: 256-261.

CHROMOSOMAL LOCATION

Genetic locus: SERF1A/SERF1B (human) mapping to 5q13.2; Serf1 (mouse) mapping to 13 D1.

SOURCE

SERF1 (K-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SERF1 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-240856 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SERF1 (K-13) is recommended for detection of SERF1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with SERF2.

Suitable for use as control antibody for SERF1 siRNA (h): sc-92056, SERF1 siRNA (m): sc-153343, SERF1 shRNA Plasmid (h): sc-92056-SH, SERF1 shRNA Plasmid (m): sc-153343-SH, SERF1 shRNA (h) Lentiviral Particles: sc-92056-V and SERF1 shRNA (m) Lentiviral Particles: sc-153343-V.

Molecular Weight of SERF1 long isoform: 12 kDa.

Molecular Weight of SERF1 short isoform: 7 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

- Falsone, S.F., et al. 2012. SERF protein is a direct modifier of amyloid fiber assembly. *Cell Rep.* 2: 358-371.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.