

SGTA (N-12): sc-240871

BACKGROUND

SGTA (small glutamine-rich tetratricopeptide repeat (TPR)-containing protein A or α), also known as SGT, hSGT or UBP (Vpu-binding protein), is a ubiquitously expressed protein that contains three TPR protein-protein interaction repeats. SGTA is believed to function as a component of the androgen receptor (AR)-chaperone-co-chaperone complex, acting as a co-chaperone involved in androgen signaling. More specifically, SGTA binds to the hinge region of the AR functions to retain the AR in the cytoplasm, thereby inhibiting androgen signaling. In addition, SGTA functions as a co-chaperone for HSP 90 and HSP 70, two proteins known to participate in apoptosis. On the basis of its role in apoptosis and androgen signaling, SGTA is a potential candidate for PCOS (polycystic ovary syndrome), a disorder characterized by androgen excess, obesity and menstrual disturbances. SGTA also interacts with the nonstructural Parvovirus protein NS1 and the HIV-1 proteins Vpu and Gag.

REFERENCES

- Kordes, E., et al. 1998. Isolation and characterization of human SGT and identification of homologues in *Saccharomyces cerevisiae* and *Caenorhabditis elegans*. *Genomics* 52: 90-94.
- Tobaben, S., et al. 2001. A trimeric protein complex functions as a synaptic chaperone machine. *Neuron* 31: 987-999.
- Angeletti, P.C., et al. 2002. Small glutamine-rich protein/viral protein U-binding protein is a novel cochaperone that affects heat shock protein 70 activity. *Cell Stress Chaperones* 7: 258-268.
- Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603419. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Tobaben, S., et al. 2003. A brain-specific isoform of small glutamine-rich tetratricopeptide repeat-containing protein binds to Hsc70 and the cysteine string protein. *J. Biol. Chem.* 278: 38376-38383.
- Liou, S.T., et al. 2005. Small glutamine-rich tetratricopeptide repeat-containing protein is composed of three structural units with distinct functions. *Arch. Biochem. Biophys.* 435: 253-263.
- Yin, H., et al. 2006. SGT, a Hsp90 β binding partner, is accumulated in the nucleus during cell apoptosis. *Biochem. Biophys. Res. Commun.* 343: 1153-1158.
- Buchanan, G., et al. 2007. Control of androgen receptor signaling in prostate cancer by the cochaperone small glutamine rich tetratricopeptide repeat containing protein α . *Cancer Res.* 67: 10087-10096.
- Goodarzi, M.O., et al. 2008. Small glutamine-rich tetratricopeptide repeat-containing protein α (SGTA), a candidate gene for polycystic ovary syndrome. *Hum. Reprod.* 23: 1214-1219.

CHROMOSOMAL LOCATION

Genetic locus: SGTA (human) mapping to 19p13.3; Sgta (mouse) mapping to 10 C1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

SGTA (N-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SGTA of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-240871 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SGTA (N-12) is recommended for detection of SGTA of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SGTA (N-12) is also recommended for detection of SGTA in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for SGTA siRNA (h): sc-97627, SGTA siRNA (m): sc-153427, SGTA shRNA Plasmid (h): sc-97627-SH, SGTA shRNA Plasmid (m): sc-153427-SH, SGTA shRNA (h) Lentiviral Particles: sc-97627-V and SGTA shRNA (m) Lentiviral Particles: sc-153427-V.

Molecular Weight of SGTA: 35 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, SH-SY5Y cell lysate: sc-3812 or K-562 whole cell lysate: sc-2203.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.



MONOS
Satisfation
Guaranteed

Try **SGTA (6A4): sc-130557** or **SGTA (F-7): sc-374031**, our highly recommended monoclonal alternatives to SGTA (N-12).