

Cytokeratin 8 (T-13): sc-241376

BACKGROUND

Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. Cytokeratin 8 expression is seen in epithelium and epithelium-derived tumors. The Cytokeratin 8 and 18 pair are normally expressed in simple epithelia, but not in stratified epithelial cells. Research indicates that squamous cell carcinomas derived from stratified epithelia show abnormal expression of Cytokeratin 8 and 18, although it is not known whether these proteins contribute to the malignant phenotype of the cells. Expression of Cytokeratin 8 and 18 in oral squamous cell carcinomas is an independent prognostic marker that indicates a poor prognosis. Cytokeratin 8 expression correlates with malignancy in leukoplakia and carcinomas of the head and neck; it is expressed in all non-small-cell lung cancers. Cytokeratin 8 has been shown to possess extracellular epitopes on tumor cells, which may represent valuable targets for therapy.

REFERENCES

1. Leube, R.E., et al. 1986. Cytokeratin expression in simple epithelia. III. Detection of mRNAs encoding human Cytokeratins nos. 8 and 18 in normal and tumor cells by hybridization with cDNA sequences *in vitro* and *in situ*. *Differentiation* 33: 69-85.
2. van der Velden, L.A., et al. 1993. Cytokeratin expression in normal and (pre)malignant head and neck epithelia: an overview. *Head Neck* 15: 133-146.
3. Silen, A., et al. 1994. Evaluation of a new tumor marker for Cytokeratin 8 and 18 fragments in healthy individuals and prostate cancer patients. *Prostate* 24: 326-332.

CHROMOSOMAL LOCATION

Genetic locus: KRT8 (human) mapping to 12q13.13; Krt8 (mouse) mapping to 15 F3.

SOURCE

Cytokeratin 8 (T-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Cytokeratin 8 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-241376 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Cytokeratin 8 (T-13) is recommended for detection of Cytokeratin 8 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Cytokeratin family members.

Suitable for use as control antibody for Cytokeratin 8 siRNA (h): sc-35156, Cytokeratin 8 siRNA (m): sc-72111, Cytokeratin 8 shRNA Plasmid (h): sc-35156-SH, Cytokeratin 8 shRNA Plasmid (m): sc-72111-SH, Cytokeratin 8 shRNA (h) Lentiviral Particles: sc-35156-V and Cytokeratin 8 shRNA (m) Lentiviral Particles: sc-72111-V.


Molecular Weight of Cytokeratin 8: 40-55 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, K-562 whole cell lysate: sc-2203 or T-47D cell lysate: sc-2293.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Cytokeratin 8 (T-13): sc-241376. Western blot analysis of Cytokeratin 8 expression in K-562 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Cytokeratin 8 (C51): sc-8020** or **Cytokeratin 8 (A-9): sc-374275**, our highly recommended monoclonal alternatives to Cytokeratin 8 (T-13). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **Cytokeratin 8 (C51): sc-8020**.