

# Ggt1 (L-15): sc-241476

## BACKGROUND

Homologous glycosyltransferase (GT) gene families catalyze the formation of glycosidic linkages. The  $\beta$ -1,3 galactosyltransferase ( $\beta$ 3GalT) gene family encodes a set of type II transmembrane glycoproteins that are catalytically diverse and use different donor substrates (UDP-galactose and UDP-N-acetylglucosamine) and different acceptor sugars (N-acetylglucosamine, galactose, N-acetylgalactosamine) to catalyze the addition of an activated monosaccharide to a terminal lactose. The  $\beta$ -1,4-galactosyltransferase ( $\beta$ 4GalT) gene family shows exclusive specificity for the donor substrate UDP-galactose. In several tissues and cell lines, GTs localize to the Golgi complex. Ggt1 (N-acetyllactosaminide  $\alpha$ -1,3-galactosyltransferase), also known as GALT or Gal, is a 394 amino acid single-pass type II membrane protein that belongs to the glycosyltransferase 6 family. Localizing to the Golgi apparatus, Ggt1 utilizes manganese as a cofactor and is involved in the transfer of galactose from UDP-galactose to acceptor molecules. Existing as three alternatively spliced isoforms, the gene encoding Ggt1 maps to mouse chromosome 2 B.

## REFERENCES

- Larsen, R.D., Rajan, V.P., Ruff, M.M., Kukowska-Latallo, J., Cummings, R.D. and Lowe, J.B. 1989. Isolation of a cDNA encoding a murine UDPgalactose:  $\beta$ -D-galactosyl-1,4-N-acetyl-D-glucosaminide  $\alpha$ -1,3-galactosyltransferase: expression cloning by gene transfer. Proc. Natl. Acad. Sci. USA 86: 8227-8231.
- Joziasse, D.H., Shaper, N.L., Kim, D., Van den Eijnden, D.H. and Shaper, J.H. 1992. Murine  $\alpha$  1,3-galactosyltransferase. A single gene locus specifies four isoforms of the enzyme by alternative splicing. J. Biol. Chem. 267: 5534-5541.
- Koike, C., Friday, R., Fung, J.J., Starzl, T.E. and Trucco, M. 2001. Comparison of the regulatory regions of the  $\alpha$  1,3galactosyltransferase gene between murine and porcine species. Transplant. Proc. 33: 710-711.
- Alexander, W.S., Viney, E.M., Zhang, J.G., Metcalf, D., Kauppi, M., Hyland, C.D., Carpinelli, M.R., Stevenson, W., Croker, B.A., Hilton, A.A., Ellis, S., Selan, C., Nandurkar, H.H., Goodnow, C.C., Kile, B.T., Nicola, N.A., Roberts, A.W. and Hilton, D.J. 2006. Thrombocytopenia and kidney disease in mice with a mutation in the C1galt1 gene. Proc. Natl. Acad. Sci. USA 103: 16442-16447.
- Xu, R., Chandrasekharan, K., Yoon, J.H., Camboni, M. and Martin, P.T. 2007. Overexpression of the cytotoxic T cell (CT) carbohydrate inhibits muscular dystrophy in the dyW mouse model of congenital muscular dystrophy 1A. Am. J. Pathol. 171: 181-199.
- Martin, P.T. 2007. Congenital muscular dystrophies involving the O-mannose pathway. Curr. Mol. Med. 7: 417-425.

## CHROMOSOMAL LOCATION

Genetic locus: Ggt1 (mouse) mapping to 2 B.

## SOURCE

Ggt1 (L-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Ggt1 of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-241476 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Ggt1 (L-15) is recommended for detection of Ggt1 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Ggt1 siRNA (m): sc-145392, Ggt1 shRNA Plasmid (m): sc-145392-SH and Ggt1 shRNA (m) Lentiviral Particles: sc-145392-V.


Molecular Weight of Ggt1 isoforms 1/2/3: 47/43/44 kDa.

Positive Controls: BC<sub>3</sub>H1 cell lysate: sc-2299.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



Ggt1 (L-15): sc-241476. Western blot analysis of Ggt1 expression in BC<sub>3</sub>H1 whole cell lysate.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.