

# Prickle4 (S-13): sc-241643

## BACKGROUND

Prickle4, also known as OEPT (overexpressed breast tumor protein), OBTP or TOMM6, is a 344 amino acid protein that contains one PET domain, two LIM zinc-binding domains, and belongs to the Prickle family. Prickle4 exists as three alternatively spliced isoforms that are expressed in prostate and breast cancers, hepatocellular carcinoma and a number of normal tissues. The gene encoding Prickle4 maps to human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Additionally, Porphyria cutanea tarda, Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder are all associated with genes that map to chromosome 6.

## REFERENCES

1. Chassin, D., Benifla, J.L., Delattre, C., Fernandez, H., Ginisty, D., Janneau, J.L., Prade, M., Contesso, G., Caillou, B. and Tournaire, M. 1994. Identification of genes overexpressed in tumors through preferential expression screening in trophoblasts. *Cancer Res.* 54: 5217-5223.
2. Brunner, H.G., van Beersum, S.E., Warman, M.L., Olsen, B.R., Ropers, H.H. and Mariman, E.C. 1994. A Stickler syndrome gene is linked to chromosome 6 near the COL11A2 gene. *Hum. Mol. Genet.* 3: 1561-1564.
3. Cesari, R., Martin, E.S., Calin, G.A., Pentimalli, F., Bichi, R., McAdams, H., Trapasso, F., Drusco, A., Shimizu, M., Masciullo, V., D'Andrilli, G., Scambia, G., Picchio, M.C., Alder, H., Godwin, A.K. and Croce, C.M. 2003. Parkin, a gene implicated in autosomal recessive juvenile parkinsonism, is a candidate tumor suppressor gene on chromosome 6q25-q27. *Proc. Natl. Acad. Sci. USA* 100: 5956-5961.
4. Teufel, A., Weinmann, A., Galle, P.R. and Lohse, A.W. 2005. Characterization of OEPT, a LIM protein. *Int. J. Mol. Med.* 15: 513-518.
5. Oh, J.H., Yang, J.O., Hahn, Y., Kim, M.R., Byun, S.S., Jeon, Y.J., Kim, J.M., Song, K.S., Noh, S.M., Kim, S., Yoo, H.S., Kim, Y.S. and Kim, N.S. 2005. Transcriptome analysis of human gastric cancer. *Mamm. Genome* 16: 942-954.
6. Bläker, H., Mechttersheimer, G., Sutter, C., Hertkorn, C., Kern, M.A., Rieker, R.J., Penzel, R., Schirmacher, P. and Kloor, M. 2008. Recurrent deletions at 6q in early age of onset non-HNPCC- and non-FAP-associated intestinal carcinomas. Evidence for a novel cancer susceptibility locus at 6q14-q22. *Genes Chromosomes Cancer* 47: 159-164.
7. Jalil, S., Grady, J.J., Lee, C. and Anderson, K.E. 2010. Associations among behavior-related susceptibility factors in porphyria cutanea tarda. *Clin. Gastroenterol. Hepatol.* 8: 297-302.

## CHROMOSOMAL LOCATION

Genetic locus: PRICKLE4 (human) mapping to 6p21.1.

## SOURCE

Prickle4 (S-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Prickle4 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-241643 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Prickle4 (S-13) is recommended for detection of Prickle4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with LMO6, Prickle1 or Prickle2.

Prickle4 (S-13) is also recommended for detection of Prickle4 in additional species, including porcine.

Suitable for use as control antibody for Prickle4 siRNA (h): sc-95577, Prickle4 shRNA Plasmid (h): sc-95577-SH and Prickle4 shRNA (h) Lentiviral Particles: sc-95577-V.

Molecular Weight of Prickle4 isoforms 1/2/3: 38/33/42 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.