

TMEM161B (G-12): sc-241707

BACKGROUND

TMEM161B (transmembrane protein 161B) is a 487 amino acid protein encoded by a gene mapping to human chromosome 5. With 181 million base pairs encoding around 1,000 genes, chromosome 5 is about 6% of human genomic DNA. It is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5 associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome. Deletion of 5q or chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

- Dixon, M.J., et al. 1991. The gene for Treacher Collins syndrome maps to the long arm of chromosome 5. *Am. J. Hum. Genet.* 49: 17-22.
- Saltman, D.L., et al. 1993. A physical map of 15 loci on human chromosome 5q23-q33 by two-color fluorescence *in situ* hybridization. *Genomics* 16: 726-732.
- Kadmon, M., et al. 2001. Duodenal adenomatosis in familial adenomatous polyposis coli. A review of the literature and results from the Heidelberg Polyposis Register. *Int. J. Colorectal Dis.* 16: 63-75.
- South, S.T., et al. 2006. A new genomic mechanism leading to cri-du-chat syndrome. *Am. J. Med. Genet. A* 140: 2714-2720.
- Aretz, S., et al. 2007. Somatic APC mosaicism: a frequent cause of familial adenomatous polyposis (FAP). *Hum. Mutat.* 28: 985-992.
- Cleaver, J.E., et al. 2007. Cockayne syndrome exhibits dysregulation of p21 and other gene products that may be independent of transcription-coupled repair. *Neuroscience* 145: 1300-1308.
- Du, H.Y., et al. 2007. Telomerase reverse transcriptase haploinsufficiency and telomere length in individuals with 5p- syndrome. *Aging Cell* 6: 689-697.
- Herry, A., et al. 2007. Redefining monosomy 5 by molecular cytogenetics in 23 patients with MDS/AML. *Eur. J. Haematol.* 78: 457-467.
- Makrantonaki, E., et al. 2007. Molecular mechanisms of skin aging: state of the art. *Ann. N.Y. Acad. Sci.* 1119: 40-50.

CHROMOSOMAL LOCATION

Genetic locus: TMEM161B (human) mapping to 5q14.3; Tmem161b (mouse) mapping to 13 C3.

SOURCE

TMEM161B (G-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TMEM161B of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-241707 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TMEM161B (G-12) is recommended for detection of TMEM161B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with AROS-29.


TMEM161B (G-12) is also recommended for detection of TMEM161B in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for TMEM161B siRNA (h): sc-91740, TMEM161B siRNA (m): sc-154390, TMEM161B shRNA Plasmid (h): sc-91740-SH, TMEM161B shRNA Plasmid (m): sc-154390-SH, TMEM161B shRNA (h) Lentiviral Particles: sc-91740-V and TMEM161B shRNA (m) Lentiviral Particles: sc-154390-V.

Molecular Weight of TMEM161B: 55 kDa.

Positive Controls: Jurkat Whole Cell Lysate: sc-2204 or K-562 Whole Cell Lysate: sc-2203.

DATA



TMEM161B (G-12): sc-241707. Western blot analysis of TMEM161B expression in K-562 (A), Jurkat (B) and MCF7 (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.