

WTIP (S-13): sc-241738

BACKGROUND

Wilms' tumor (WT) is an embryonal malignancy of the kidney that affects 1 in 10,000 infants and is observed in both sporadic and inherited forms. The Wilms' tumor locus has been mapped at chromosome 11p13.11 as a tumor suppressor gene which encodes a DNA binding protein with four zinc fingers and a glutamine-proline rich amino-terminus. WTIP (Wilms tumor 1 interacting protein) is a 654 amino acid LIM domain protein that belongs to the zyxin/ajuba family, interacts with WT1 (Wilms tumor 1) and is thought to play a role in slit diaphragm protein assembly. Acting as a transcription regulator, WTIP shuttles between nucleus and adhesion structures following podocyte injury to repress WT1-dependent transcription regulation. WTIP contains three LIM zinc-binding domains and is encoded by a gene that maps to human chromosome 19q13.11.

REFERENCES

- Weissman, B.E., et al. 1987. Introduction of a normal human chromosome 11 into a Wilms' tumor cell line controls its tumorigenic expression. *Science* 236: 175-180.
- Morris, J.F., et al. 1991. Characterization of the zinc finger protein encoded by the WT1 Wilms' tumor locus. *Oncogene* 6: 2339-2348.
- Srichai, M.B., et al. 2004. A WT1 co-regulator controls podocyte phenotype by shuttling between adhesion structures and nucleus. *J. Biol. Chem.* 279: 14398-14408.
- Yang, L., et al. 2007. A tumor suppressor and oncogene: the WT1 story. *Leukemia* 21: 868-876.
- Langer, E.M., et al. 2008. Ajuba LIM proteins are snail/slug corepressors required for neural crest development in *Xenopus*. *Dev. Cell* 14: 424-436.
- van Wijk, N.V., et al. 2009. The LIM domain protein Wtip interacts with the receptor tyrosine kinase Ror2 and inhibits canonical Wnt signalling. *Biochem. Biophys. Res. Commun.* 390: 211-216.
- Kim, J.H., et al. 2010. Podocyte injury induces nuclear translocation of WTIP via microtubule-dependent transport. *J. Biol. Chem.* 285: 9995-10004.
- Hou, Z., et al. 2010. LIM protein Ajuba functions as a nuclear receptor corepressor and negatively regulates retinoic acid signaling. *Proc. Natl. Acad. Sci. USA* 107: 2938-2943.

CHROMOSOMAL LOCATION

Genetic locus: WTIP (human) mapping to 19q13.11; Wtip (mouse) mapping to 7 B1.

SOURCE

WTIP (S-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of WTIP of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-241738 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

WTIP (S-13) is recommended for detection of WTIP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

WTIP (S-13) is also recommended for detection of WTIP in additional species, including canine and bovine.

Suitable for use as control antibody for WTIP siRNA (m): sc-155365, WTIP shRNA Plasmid (m): sc-155365-SH and WTIP shRNA (m) Lentiviral Particles: sc-155365-V.


Molecular Weight of WTIP: 67 kDa.

Positive Controls: WTIP transfected CHO whole cell lysate or WiDR cell lysate: sc-24779.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



WTIP (S-13): sc-241738. Western blot analysis of WTIP expression in non-transfected CHO (A), human WTIP transfected CHO (B) and WiDR (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.