

# ASPDH (T-17): sc-241845

## BACKGROUND

ASPDH (aspartate dehydrogenase domain containing), also known as putative L-aspartate dehydrogenase, is a 283 amino acid belonging to the L-aspartate dehydrogenase family. ASPDH participates in NADP and NADPH binding, as well as aspartate dehydrogenase and oxidoreductase activities. ASPDH catalyzes NAD and NADP-dependent dehydrogenation of L-aspartate to iminoaspartate, resulting in an unstable iminoaspartate product, which can decompose to oxaloacetate and ammonia. Existing as two alternatively spliced isoforms, ASPDH is encoded by a gene that maps to human chromosome 19q13.33. Chromosome 19 makes up over 2% of the human genome and contains approximately 63 million bases, which encode over 1,400 genes. Recognized for having the greatest gene density of all human chromosomes, chromosome 19 is linked to Peutz-Jeghers syndrome, spinocerebellar ataxia type 6, the stroke disorder CADASIL, hypercholesterolemia and Insulin-dependent diabetes. Translocation of chromosomes 19 and 14 may be related to lymphoproliferative disorders.

## REFERENCES

- Zimmermann, W., et al. 1988. Chromosomal localization of the carcinoembryonic antigen gene family and differential expression in various tumors. *Cancer Res.* 48: 2550-2554.
- LaPoint, S.F., et al. 2000. Cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL). *Adv. Anat. Pathol.* 7: 307-321.
- Trettel, F., et al. 2000. A fine physical map of the CACNA1A gene region on 19p13.1-p13.2 chromosome. *Gene* 241: 45-50.
- Buchet-Poyau, K., et al. 2002. Search for the second Peutz-Jeghers syndrome locus: exclusion of the STK13, PRKCG, KLK10, and PSCD2 genes on chromosome 19 and the STK11IP gene on chromosome 2. *Cytogenet. Genome Res.* 97: 171-178.
- Moodie, S.J., et al. 2002. Analysis of candidate genes on chromosome 19 in coeliac disease: an association study of the KIR and LILR gene clusters. *Eur. J. Immunogenet.* 29: 287-291.
- Grimwood, J., et al. 2004. The DNA sequence and biology of human chromosome 19. *Nature* 428: 529-535.
- Parham, P. 2005. Immunogenetics of killer cell immunoglobulin-like receptors. *Mol. Immunol.* 42: 459-462.

## CHROMOSOMAL LOCATION

Genetic locus: ASPDH (human) mapping to 19q13.33; Aspdh (mouse) mapping to 7 B4.

## SOURCE

ASPDH (T-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ASPDH of human origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-241845 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

ASPDH (T-17) is recommended for detection of ASPDH of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ASPDH (T-17) is also recommended for detection of ASPDH in additional species, including porcine and canine.

Suitable for use as control antibody for ASPDH siRNA (h): sc-97089, ASPDH siRNA (m): sc-108114, ASPDH shRNA Plasmid (h): sc-97089-SH, ASPDH shRNA Plasmid (m): sc-108114-SH, ASPDH shRNA (h) Lentiviral Particles: sc-97089-V and ASPDH shRNA (m) Lentiviral Particles: sc-108114-V.


Molecular Weight of ASPDH: 30 kDa.

Positive Controls: SK-MEL-24 whole cell lysate: sc-364259.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



ASPDH (T-17): sc-241845. Western blot analysis of ASPDH expression in SK-MEL-24 whole cell lysate.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.