

NDNF (P-20): sc-242196

BACKGROUND

NDNF (neuron-derived neurotrophic factor), also known as NORD or C4orf31, is a 568 amino acid secreted protein involved in cell adhesion and matrix assembly. NDNF contains two fibronectin type-III domains, binds chondroitin sulfate and heparin, and becomes O-glycosylated following translation. The gene encoding NDNF maps to human chromosome 4, which represents approximately 6% of the human genome and contains nearly 900 genes. Notably, the Huntingtin gene, which is found to encode an expanded glutamine tract in cases of Huntington's disease, is on chromosome 4. FGFR-3 is also encoded by a gene that maps to human chromosome 4 and has been associated with thanatophoric dwarfism, achondroplasia, Muenke syndrome and bladder cancer. Chromosome 4 is also tied to Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

REFERENCES

1. Bonaventure, J., et al. 1996. Common mutations in the fibroblast growth factor receptor 3 (FGFR 3) gene account for achondroplasia, hypochondroplasia, and thanatophoric dwarfism. *Am. J. Med. Genet.* 63: 148-154.
2. Kalchman, M.A., et al. 1996. Huntingtin is ubiquitinated and interacts with a specific ubiquitin-conjugating enzyme. *J. Biol. Chem.* 271: 19385-19394.
3. Singhrao, S.K., et al. 1998. Huntingtin protein colocalizes with lesions of neurodegenerative diseases: an investigation in Huntington's, Alzheimer's, and Pick's diseases. *Exp. Neurol.* 150: 213-222.
4. Krakow, D., et al. 2000. Exclusion of the Ellis-van Creveld region on chromosome 4p16 in some families with asphyxiating thoracic dystrophy and short-rib polydactyly syndromes. *Eur. J. Hum. Genet.* 8: 645-648.
5. Sommardahl, C., et al. 2001. Phenotypic variations of orpk mutation and chromosomal localization of modifiers influencing kidney phenotype. *Physiol. Genomics* 7: 127-134.
6. Dobson, C.M., et al. 2002. Identification of the gene responsible for the cblA complementation group of vitamin B12-responsive methylmalonic acidemia based on analysis of prokaryotic gene arrangements. *Proc. Natl. Acad. Sci. USA* 99: 15554-15559.
7. Utine, G.E., et al. 2006. Mosaicism for terminal deletion of 4q. *Genet. Couns.* 17: 205-209.

CHROMOSOMAL LOCATION

Genetic locus: NDNF (human) mapping to 4q27; Ndnf (mouse) mapping to 6 C1.

SOURCE

NDNF (P-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of NDNF of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-242196 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NDNF (P-20) is recommended for detection of NDNF of human and mouse origin and the corresponding rat homolog of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NDNF (P-20) is also recommended for detection of NDNF in additional species, including equine, canine, porcine and avian.

Suitable for use as control antibody for NDNF siRNA (h): sc-88858, NDNF siRNA (m): sc-140706, NDNF shRNA Plasmid (h): sc-88858-SH, NDNF shRNA Plasmid (m): sc-140706-SH, NDNF shRNA (h) Lentiviral Particles: sc-88858-V and NDNF shRNA (m) Lentiviral Particles: sc-140706-V.


Molecular Weight of NDNF: 65 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204 or K-562 whole cell lysate: sc-2203.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



NDNF1 (P-20): sc-242196. Western blot analysis of NDNF expression in Jurkat (A) and K-562 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.