

## LENG4 (S-19): sc-243243

### BACKGROUND

The membrane-bound O-acyltransferase family is a group of integral membrane proteins with acyltransferase activity. LENG4 (membrane bound O-acyltransferase domain containing 7), also known as lysophospholipid acyltransferase 7, leukocyte receptor cluster member 4, MBOAT7, BB1, LPLAT 7, LRC4 or OACT7, is a 472 amino acid multi-pass membrane protein involved in lipid and phospholipid metabolism. LENG4 belongs to the membrane-bound acyltransferase family and participates in the Land cycle by catalyzing reacylation of phospholipids. LENG4 is responsible for the conversion of lysophosphatidylinositol into phosphatidylinositol, and prefers arachidonoyl-CoA as its acyl donor. Overexpressed in metastatic breast and bladder carcinomas, LENG4 is down-regulated by gamma-interferon. Three LENG4 isoforms exist due to alternative splicing events, and LENG4 is encoded by a gene mapping to human chromosome 19q13.42.

### REFERENCES

1. Fukunaga-Johnson, N., Lee, S.W., Liebert, M. and Grossman, H.B. 1996. Molecular analysis of a gene, BB1, overexpressed in bladder and breast carcinoma. *Anticancer Res.* 16: 1085-1090.
2. Wende, H., Volz, A. and Ziegler, A. 2000. Extensive gene duplications and a large inversion characterize the human leukocyte receptor cluster. *Immunogenetics* 51: 703-713.
3. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 606048. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Matsuda, S., Inoue, T., Lee, H.C., Kono, N., Tanaka, F., Gengyo-Ando, K., Mitani, S. and Arai, H. 2008. Member of the membrane-bound O-acyltransferase (MBOAT) family encodes a lysophospholipid acyltransferase with broad substrate specificity. *Genes Cells* 13: 879-888.
5. Gijón, M.A., Riekhof, W.R., Zarini, S., Murphy, R.C. and Voelker, D.R. 2008. Lysophospholipid acyltransferases and arachidonate recycling in human neutrophils. *J. Biol. Chem.* 283: 30235-30245.
6. Shindou, H., Hishikawa, D., Harayama, T., Yuki, K. and Shimizu, T. 2009. Recent progress on acyl CoA: lysophospholipid acyltransferase research. *J. Lipid Res.* 50: S46-S51.

### CHROMOSOMAL LOCATION

Genetic locus: MBOAT7 (human) mapping to 19q13.42; Mboat7 (mouse) mapping to 7 A1.

### SOURCE

LENG4 (S-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of LENG4 of human origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-243243 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

LENG4 (S-19) is recommended for detection of LENG4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with LENG1, LENG8 or LENG9.

LENG4 (S-19) is also recommended for detection of LENG4 in additional species, including equine and canine.

Suitable for use as control antibody for LENG4 siRNA (h): sc-97194, LENG4 siRNA (m): sc-146703, LENG4 shRNA Plasmid (h): sc-97194-SH, LENG4 shRNA Plasmid (m): sc-146703-SH, LENG4 shRNA (h) Lentiviral Particles: sc-97194-V and LENG4 shRNA (m) Lentiviral Particles: sc-146703-V.

Molecular Weight of LENG4 isoforms: 55/45/38 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.