

LHFPL2 (G-15): sc-243259

BACKGROUND

The development of lipomas, benign tumors composed of fatty tissues, has been linked to breakpoints in the HMGI-C gene. LHFP (lipoma HMGIC fusion partner) is a multi-pass membrane protein that acts as a fusion partner with HMGI-C in a lipoma with the translocation t(12;13)(q13-q15;q12). An LHFP family member, LHFPL2 (lipoma HMGIC fusion partner-like 2 protein) is a 228 amino acid multi-pass membrane protein that is expressed in most tissues except brain and peripheral blood leukocytes. The gene encoding LHFPL2 maps to human chromosome 5q14.1, which is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5 associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome. Deletion of 5q or chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

- Ishwad, C.S., et al. 1997. The high mobility group I-C gene (HMGI-C): polymorphism and genetic localization. *Hum. Genet.* 99: 103-105.
- Petit, M.M., et al. 1999. LHFP, a novel translocation partner gene of HMGI-C in a lipoma, is a member of a new family of LHFP-like genes. *Genomics* 57: 438-441.
- Rogalla, P., et al. 2002. Absence of HMGI-C-LHFP fusion in pulmonary chondroid hamartomas with aberrations involving chromosomal regions 12q13 through 15 and 13q12 through q14. *Cancer Genet. Cytogenet.* 133: 90-93.
- Rauch, A., et al. 2007. Chromosome 5q subtelomeric deletion syndrome. *Am. J. Med. Genet. C Semin. Med. Genet.* 145C: 372-376.
- Shadduck, R.K., et al. 2007. Recent advances in myelodysplastic syndromes. *Exp. Hematol.* 35: 137-143.
- Kristoffersen, K.E. 2008. Speech and language development in cri du chat syndrome: a critical review. *Clin. Linguist. Phon.* 22: 443-457.
- Valent, P. 2008. Revealing the pathogenesis of the 5q- syndrome. *Eur. J. Clin. Invest.* 38: 539-540.
- Buysse, K., et al. 2008. Mapping of 5q35 chromosomal rearrangements within a genomically unstable region. *J. Med. Genet.* 45: 672-678.
- Azman, B.Z., et al. 2008. Two cases of deletion 5p syndrome: one with paternal involvement and another with atypical presentation. *Singapore Med. J.* 49: e98-e100.

CHROMOSOMAL LOCATION

Genetic locus: LHFPL2 (human) mapping to 5q14.1.

SOURCE

LHFPL2 (G-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of LHFPL2 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-243259 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LHFPL2 (G-15) is recommended for detection of LHFPL2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

LHFPL2 (G-15) is also recommended for detection of LHFPL2 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for LHFPL2 siRNA (h): sc-91990, LHFPL2 shRNA Plasmid (h): sc-91990-SH and LHFPL2 shRNA (h) Lentiviral Particles: sc-91990-V.

Molecular Weight of LHFPL2: 24 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotting A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.