

LPAAT- η (N-17): sc-243309

BACKGROUND

Phosphatidic acid and lysophosphatidic acid are phospholipids involved in lipid biosynthesis and signal transduction. LPAAT- η , also known as lysophospholipid acyltransferase LPCAT4, AGPAT7 (1-acylglycerol-3-phosphate O-acyltransferase 7), AYTL3 (acyltransferase-like 3) or LPEAT2 (lysophosphatidylethanolamine acyltransferase 2), is a 524 amino acid protein belonging to the 1-acyl-sn-glycerol-3-phosphate acyltransferase family. LPAAT- η displays acyl-CoA-dependent lysophospholipid acyltransferase activity, with lysophospholipids as its substrates. For example, LPAAT- η converts lysophosphatidylethanolamine to phosphatidylethanolamine and lysophosphatidylcholine to phosphatidylcholine, respectively. In contrast, LPAAT- η has no lysophosphatidylinositol, glycerol-3-phosphate, diacylglycerol or lysophosphatidic acid acyltransferase activity. LPAAT- η also prefers long chain acyl-CoAs (C16, C18) as acyl donors. Localized to the endoplasmic reticulum membrane, LPAAT- η is widely expressed with predominant levels in brain. Two isoforms of LPAAT- η are produced by alternative splicing events.

REFERENCES

1. Siegel, H.I. and Rosenblatt, J.S. 1975. Latency and duration of estrogen induction of maternal behavior in hysterectomized-ovariectomized virgin rats: effects of pup stimulation. *Physiol. Behav.* 14: 473-476.
2. West, J., Tompkins, C.K., Balantac, N., Nudelman, E., Meengs, B., White, T., Bursten, S., Coleman, J., Kumar, A., Singer, J.W. and Leung, D.W. 1997. Cloning and expression of two human lysophosphatidic acid acyltransferase cDNAs that enhance cytokine-induced signaling responses in cells. *DNA Cell Biol.* 16: 691-701.
3. Aguado, B. and Campbell, R.D. 1998. Characterization of a human lysophosphatidic acid acyltransferase that is encoded by a gene located in the class III region of the human major histocompatibility complex. *J. Biol. Chem.* 273: 4096-4105.
4. Bursten, S.L. 1998. Interaction of lipopolysaccharide with a mammalian lyso-phosphatidate acyltransferase (LPAAT) transfected into *E. coli*, and effect of lisofylline on LPAAT transfected into mammalian cells. *Prog. Clin. Biol. Res.* 397: 345-356.
5. Eberhardt, C., Gray, P.W. and Tjoelker, L.W. 1999. cDNA cloning, expression and chromosomal localization of two human lysophosphatidic acid acyltransferases. *Adv. Exp. Med. Biol.* 469: 351-356.
6. Ye, G.M., Chen, C., Huang, S., Han, D.D., Guo, J.H., Wan, B. and Yu, L. 2005. Cloning and characterization a novel human 1-acyl-sn-glycerol-3-phosphate acyltransferase gene AGPAT7. *DNA Seq.* 16: 386-390.
7. Cao, J., Shan, D., Revett, T., Li, D., Wu, L., Liu, W., Tobin, J.F. and Gimeno, R.E. 2008. Molecular identification of a novel mammalian brain isoform of acyl-CoA:lysophospholipid acyltransferase with prominent ethanolamine lysophospholipid acylating activity, LPEAT2. *J. Biol. Chem.* 283: 19049-19057.
8. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 612039. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/612039>

CHROMOSOMAL LOCATION

Genetic locus: LPCAT4 (human) mapping to 15q14; Lpcat4 (mouse) mapping to 2 E3.

SOURCE

LPAAT- η (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of LPAAT- η of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-243309 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LPAAT- η (N-17) is recommended for detection of LPAAT- η of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other LPAAT family members.

LPAAT- η (N-17) is also recommended for detection of LPAAT- η in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for LPAAT- η siRNA (h): sc-90140, LPAAT- η siRNA (m): sc-149019, LPAAT- η shRNA Plasmid (h): sc-90140-SH, LPAAT- η shRNA Plasmid (m): sc-149019-SH, LPAAT- η shRNA (h) Lentiviral Particles: sc-90140-V and LPAAT- η shRNA (m) Lentiviral Particles: sc-149019-V.

Molecular Weight of LPAAT- η isoforms 1/2: 57/49 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.